

# SPACES OF COINVARIANTS AND FUSION PRODUCT II. $\widehat{\mathfrak{sl}}_2$ CHARACTER FORMULAS IN TERMS OF KOSTKA POLYNOMIALS

B. FEIGIN, M. JIMBO, R. KEDEM, S. LOKTEV, T. MIWA

**ABSTRACT.** In this paper, we continue our study of the Hilbert polynomials of coinvariants begun in our previous work [FJKLM] (paper I). We describe the  $\mathfrak{sl}_n$  fusion products for symmetric tensor representations following the method of [FF], and show that their Hilbert polynomials are  $A_{n-1}$ -supernomials. We identify the fusion product of arbitrary irreducible  $\mathfrak{sl}_n$ -modules with the fusion product of their restriction to  $\mathfrak{sl}_{n-1}$ . Then using the equivalence theorem from paper I and the results above for  $\mathfrak{sl}_3$  we give a fermionic formula for the Hilbert polynomials of a class of  $\widehat{\mathfrak{sl}}_2$  coinvariants in terms of the level-restricted Kostka polynomials. The coinvariants under consideration are a generalization of the coinvariants studied in [FKLMM]. Our formula differs from the fermionic formula established in [FKLMM] and implies the alternating sum formula conjectured in [FL] for this case.

## 1. INTRODUCTION

Combinatorics and representation theory share a long history of mutual interaction. Calculation of dimensions, multiplicities, etc., provides interesting problems for combinatorics, and combinatorics gives a natural language and technical machinery for solving them. In many cases, representation theory provides an interpretation of combinatorial identities, or creates new identities.

The Kostka-Foulkes polynomial  $K_{\lambda\mu}(q)$ , the transition matrix between Hall Littlewood and Schur functions, is one of the main players in this interaction. In the original version, where  $\lambda$  and  $\mu$  are partitions, the Kostka number  $K_{\lambda\mu}(1)$  is the cardinality of the set of Young tableaux  $\mathcal{T}(\lambda, \mu)$  of shape  $\lambda$  and weight  $\mu$ . It gives the multiplicity of the irreducible  $\mathfrak{sl}_n$ -module of highest weight  $\lambda$  in the tensor product of the  $\mu_i$ -th symmetric tensor representations ( $1 \leq i \leq N$ ).

In combinatorics, introduction of  $q$ -polynomials such as  $K_{\lambda\mu}(q)$  is equivalent to finding a good statistic on combinatorial objects. In [LS], it was proved that the charge statistic on  $\mathcal{T}(\lambda, \mu)$  gives the Kostka polynomial.

In recent years, the interaction between combinatorics and representation theory has been enhanced by integrable models in statistical mechanics and

quantum field theory. This is because physics provides the equivalent of a charge statistic.

For the Kostka polynomial, [KR] found a charge preserving bijection between Young tableaux and rigged configurations. The latter appear naturally in the context of counting the Bethe Ansatz states of the XXX Hamiltonian of one-dimensional quantum spin chain. The meaning of the charge statistic for the Bethe states was later clarified in [KKMM], where it was found that in the conformal limit, the scaled partition function in the infinite size limit, corresponding to the Bethe states, in many integrable models gives characters in conformal field theory (CFT). The bijection of [KR] led to new combinatorial identities which express the Kostka polynomials as sums of products of  $q$  binomial coefficients. Such a formula was called a fermionic formula in [KKMM].

Although a variety of fermionic formulas and related identities have been obtained by combinatorial methods, the meaning from representation theory was not given in general. In [FJKLM] we clarified some aspect of this connection between filtered spaces of conformal blocks and the Kostka polynomial from representation theory of affine Lie algebra  $\widehat{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}K$ . Namely, we defined a natural filtration on the dual space of conformal blocks, and showed that the Hilbert polynomial of the associated graded space gives rise to the Kostka polynomial and its generalizations. Let us recall this construction.

Let  $\mathcal{Z}$  be a set of distinct complex numbers. To each  $z_i$  we associate a left ideal  $\mathfrak{X}_i \subset U(\mathfrak{g})$ . We fix a level  $k$ , and consider the finite-dimensional cyclic  $\mathfrak{g}$ -module

$$\pi^{(k)}(\mathfrak{X}_i) = \bigoplus_{\mu \in P_+^{(k)}} \left( \pi_\mu^{\mathfrak{X}_i} \right)^* \otimes \pi_\mu.$$

Here  $\pi_\mu$  is the irreducible  $\mathfrak{g}$ -module with highest weight  $\mu$ , and  $(\pi_\mu^{\mathfrak{X}_i})^*$  is the dual space of  $\mathfrak{X}_i$  invariants in  $\pi_\mu$ . The highest weight  $\mu$  is restricted by the condition  $\langle \mu, \theta^\vee \rangle \leq k$  where  $\theta^\vee$  is the maximal coroot.

Let  $L_i$  be a level  $k$  integrable (in general, reducible)  $\widehat{\mathfrak{g}}$ -module localized at  $z_i$ , which has  $\pi^{(k)}(\mathfrak{X}_i)$  as the degree zero part. Let us consider the space of conformal coinvariants

$$\begin{aligned} & \langle L_0, L_1, \dots, L_N \rangle_{\infty, z_1, \dots, z_N}^{\mathbb{C}P^1} \\ (1.1) \quad &= L_0 \boxtimes L_1 \boxtimes \dots \boxtimes L_N / \mathfrak{g}^{\mathbb{C}P^1 \setminus \{\infty, z_1, \dots, z_N\}}(L_0 \boxtimes L_1 \boxtimes \dots \boxtimes L_N), \end{aligned}$$

where  $L_0 = L_\lambda^{(k)}(\infty)$  is the level  $k$  irreducible highest weight  $\widehat{\mathfrak{g}}$ -module with highest weight  $\lambda$  ‘placed at  $\infty$ ’, whereas  $L_i$  are placed at  $z_i$  (see [FJKLM] for the precise meaning of these terms). We fix a cyclic vector  $v_i \in \pi^{(k)}(\mathfrak{X}_i) \subset L_i$  to be the canonical element corresponding to the inclusions  $\pi_\mu^{\mathfrak{X}_i} \rightarrow \pi_\mu$  ( $\mu \in P_+^{(k)}$ ). Set  $X_i = S(\mathfrak{X}_i) + B_1$ , where  $B_1 = \mathfrak{g} \otimes t\mathbb{C}[t]U(\mathfrak{g}[t])$ , and  $S$  is the antipode. Note that the left ideal  $S(X_i)$  annihilates  $v_i$ . In the following, for a module  $W$  over an algebra  $A$  and a right ideal  $Y \subset A$ , we abbreviate the space of coinvariants  $W/YW$  to  $W/Y$ .

In [FJKLM] we introduced the notion of the fusion right ideal  $X_1 \otimes \cdots \otimes X_N(\mathcal{Z})$ , and proved two canonical isomorphisms

$$(1.2) \quad L_\lambda^{(k)}(\infty)/X_1 \otimes \cdots \otimes X_N(\mathcal{Z}) \rightarrow \langle L_0, L_1, \dots, L_N \rangle_{\infty, z_1, \dots, z_N}^{\mathbf{CP}^1}$$

and

$$(1.3) \quad \mathcal{F}_{\mathcal{Z}} \left( \pi^{(k)}(\mathfrak{X}_1), \dots, \pi^{(k)}(\mathfrak{X}_N) \right) / S \left( I_\lambda^{(k)} \right) \rightarrow \langle L_0, L_1, \dots, L_N \rangle_{\infty, z_1, \dots, z_N}^{\mathbf{CP}^1}.$$

Here  $\mathcal{F}_{\mathcal{Z}} \left( \pi^{(k)}(\mathfrak{X}_1), \dots, \pi^{(k)}(\mathfrak{X}_N) \right)$  is the filtered tensor product (in the sense of [FL]) of  $\pi^{(k)}(\mathfrak{X}_i)$  with the cyclic vector  $v_1 \otimes \cdots \otimes v_N$ , and  $I_\lambda^{(k)} \subset U(\mathfrak{g}[t])$  is a left ideal which annihilates the highest weight vector  $v_\lambda \in L_\lambda^{(k)}(\infty)$  (see [FJKLM] for the precise definition).

The logical steps for the proof of these isomorphisms (1.2) and (1.3), given in [FJKLM], was as follows. A standard argument shows that there are canonical surjections. The dimension of the space of conformal coinvariants (i.e., the right hand sides of (1.2) and (1.3)) is given by the Verlinde rule. We showed that the dimension of the space in the left hand side of (1.2) obeys the same Verlinde rule. Then, we proved the equivalence theorem of filtered vector spaces

$$(1.4) \quad L_\lambda^{(k)}(\infty)/X_1 \otimes \cdots \otimes X_N(\mathcal{Z}) \simeq \mathcal{F}_{\mathcal{Z}} \left( \pi^{(k)}(\mathfrak{X}_1), \dots, \pi^{(k)}(\mathfrak{X}_N) \right) / S \left( I_\lambda^{(k)} \right).$$

In [FJKLM], we studied the case where  $\mathfrak{g} = \mathfrak{sl}_2$  and the representations  $L_i$  are irreducible. In that special case, we proved that the associated graded space of (1.2)–(1.3) is isomorphic to the space of coinvariants

$$(1.5) \quad \pi^{(k)}(\mathfrak{X}_1) * \cdots * \pi^{(k)}(\mathfrak{X}_N)(\mathcal{Z}) / S \left( I_\lambda^{(k)} \right),$$

where  $\pi^{(k)}(\mathfrak{X}_1) * \cdots * \pi^{(k)}(\mathfrak{X}_N)(\mathcal{Z})$  signifies the fusion product, i.e., the associated graded space of  $\mathcal{F}_{\mathcal{Z}} \left( \pi^{(k)}(\mathfrak{X}_1), \dots, \pi^{(k)}(\mathfrak{X}_N) \right)$ . We also gave an explicit fermionic formula for the Hilbert polynomial of (1.5). In what follows we refer to the latter as the ‘character’.

The aim of the present article is to obtain similar results for a class of reducible representations  $L_i$ . Let us describe the main points. Denote by  $e, f, h$  the standard generators of  $\mathfrak{g} = \mathfrak{sl}_2$ . In [FJKLM], we studied the space of coinvariants (1.1) by taking  $\mathfrak{X}_i$  such that  $\pi^{(k)}(\mathfrak{X}_i)$  is an irreducible  $(l+1)$ -dimensional module  $\pi_l$  of  $\mathfrak{g}$ . Here we choose  $\mathfrak{X}_i$  of the form

$$U(\mathfrak{g})e^{m_i+1} + U(\mathfrak{g})f \quad (1 \leq m_i \leq k).$$

In this case we have

$$(1.6) \quad \pi^{(k)}(\mathfrak{X}_i) \simeq \oplus_{l=0}^{m_i} \pi_l,$$

which is a reducible, cyclic module over  $\mathfrak{g}$ . We choose the cyclic vector to be the sum  $\sum_{l=0}^{m_i} v_l$  of the lowest weight vectors  $v_l \in \pi_l$ . In this setting, we compute the character of the associated graded space of the space of conformal coinvariants

$$(1.7) \quad \text{gr} \langle L_0, L_1, \dots, L_N \rangle_{\infty, z_1, \dots, z_N}^{\mathbf{CP}^1}.$$

For that purpose, we take the following approach. The direct sum (1.6) can be viewed as the  $m_i$ -th symmetric tensor representation  $\tilde{\pi}_{m_i}$  of  $\mathfrak{sl}_3$  restricted to the subalgebra  $\mathfrak{sl}_2$ . We will show that the filtered tensor product appearing in (1.4) for the reducible  $\mathfrak{sl}_2$ -modules (1.6) is isomorphic to that for the irreducible  $\mathfrak{sl}_3$ -modules  $\tilde{\pi}_{m_i}$  with the cyclic vector  $v_{m_i}$ . This observation enables us to compute the right hand side of (1.4). As a result, we obtain a fermionic formula involving level-restricted Kostka polynomials for  $\mathfrak{sl}_2$  and  $q$ -binomial symbols. At  $q = 1$ , this fermionic formula reduces to the known formula for the dimension of the space of conformal blocks. This establishes the isomorphism between (1.7) and the space of coinvariants of the fusion product (1.5). The fermionic formula obtained above also settles a conjecture of [FL] in the special case  $\mathfrak{g} = \mathfrak{sl}_2$  (see Theorem 3.8). Our technical machinery from representation theory is not yet strong enough to prove the conjecture in general. We hope for a further progress in this direction.

In the case  $m_1 = \cdots = m_N = k$  with  $z_1, \dots, z_N$  being distinct, the left hand side of (1.4) reduces to the quotient of  $L_\lambda^{(k)}(\infty)$  by the right ideal generated by

$$e \otimes t^i \prod_{j=1}^N (t - z_j), \quad f \otimes t^i \quad (i \geq 0).$$

In [FKLMM], it was shown that its associated graded space coincides with the quotient of  $L_\lambda^{(k)}(\infty)$  by the ideal of the above form with  $z_1 = \cdots = z_N = 0$ . Moreover a monomial basis of that space was constructed. The method of [FKLMM] is based on the loop Heisenberg algebra contained in  $\widehat{\mathfrak{sl}_3}$ . As we mentioned already, the fermionic formula which naturally arises in this method is different from the one given in this paper. Though the two approaches are different, one could say that there is a common flavor in that both make use of the  $\mathfrak{sl}_3$  structure. It would be an interesting problem to derive analogous results in the present setting as well.

The plan of the paper is as follows. In Section 2, we extend some result of [FF] on the fusion product in the case of  $\mathfrak{sl}_n$ -modules. In particular, we obtain the equality of the Hilbert polynomial of fusion product for the symmetric tensor representations with the corresponding case of the  $A_{n-1}$  supernomials. In Section 3, we apply the result of Section 2 for the  $\mathfrak{sl}_3$  case in the computation of the space of coinvariants in the  $\mathfrak{sl}_2$  fusion product. Appendix is given for connecting filtered tensor products of  $\mathfrak{sl}_{n-1}$  and  $\mathfrak{sl}_n$ .

## 2. FUSION PRODUCT OF MODULES OVER ABELIAN LIE ALGEBRAS

In this section, we extend some of the results of [FF] to the  $\mathfrak{sl}_n$  setting for the case of symmetric tensor representations. Namely we consider the fusion product of symmetric tensor representations of  $\mathfrak{sl}_n$ , and give an explicit characterization of the annihilating ideals. As in [FF], we derive short exact sequences of these modules and obtain a decomposition of  $\mathfrak{sl}_n$  fusion products into subquotients, each being isomorphic to an  $\mathfrak{sl}_{n-1}$  fusion product. As a

corollary, we identify their characters with the  $q$ -supernomial coefficients. We also give a recursive construction of a basis of fusion products in terms of abelian currents.

**2.1. Symmetric tensors for  $\mathfrak{sl}_n$ .** We denote the generators of  $\mathfrak{sl}_n$  by  $e_{ab}, e_{ba}$  and  $h_{ab} = e_{aa} - e_{bb}$  ( $1 \leq a < b \leq n$ ), where  $e_{ab} = (\delta_{ia}\delta_{jb})_{1 \leq i, j \leq n}$  in the defining representation. We fix a standard embedding  $\mathfrak{sl}_2 \subset \cdots \subset \mathfrak{sl}_{n-1} \subset \mathfrak{sl}_n$ , regarding  $\mathfrak{sl}_i$  as the subalgebra generated by  $e_{ab}$  ( $n-i+1 \leq a \neq b \leq n$ ). In general, for a Lie algebra  $\mathfrak{g}$  we set  $\mathfrak{g}[t] = \mathfrak{g} \otimes \mathbb{C}[t]$  and write

$$x[i] = x \otimes t^i \quad (x \in \mathfrak{g}, i \geq 0).$$

Fix a set of distinct complex numbers

$$(2.1) \quad \mathcal{Z} = (z_1, \dots, z_N), \quad z_i \neq z_j \ (i \neq j).$$

Consider the  $k$ -th symmetric tensor representation  $W_k = S^k(\mathbb{C}^n)$  with lowest weight vector  $w_k$ :  $e_{ab}w_k = 0$  for  $a > b$ . Our main concern is the fusion product [FL]

$$(2.2) \quad W_{k_1} * \cdots * W_{k_N}(\mathcal{Z}),$$

defined by choosing  $w_{k_p}$  as the cyclic vector for each  $W_{k_p}$ .

A particularly simple feature of the symmetric tensors is that they can be described in terms of the abelian subalgebra

$$\mathfrak{a} = \mathbb{C}x_1 \oplus \cdots \oplus \mathbb{C}x_{n-1} \subset \mathfrak{sl}_n, \quad x_a = e_{n-a, n}.$$

Indeed,  $W_k$  is generated by  $w_k$  also as an  $\mathfrak{a}$ -module. Moreover, since  $w_k$  is annihilated by all generators  $e_{ab}$  ( $a \neq b$ ) except for  $x_1, \dots, x_{n-1}$ , the fusion product as  $\mathfrak{a}[t]$ -module is the same as the restriction to  $\mathfrak{a}[t]$  of the fusion product (2.2) as  $\mathfrak{sl}_n[t]$ -module. For that reason we will henceforth forget the  $\mathfrak{sl}_n$ -structure and focus attention to the  $\mathfrak{a}$ -structure.

The universal enveloping algebra  $U(\mathfrak{a})$  is nothing but the polynomial ring  $\mathbb{C}[x_1, \dots, x_{n-1}]$ . The annihilating ideal

$$\text{Ann}(W_k) = \{x \in \mathbb{C}[x_1, \dots, x_{n-1}] \mid xw_k = 0\}$$

is generated by the elements  $x_1^{\nu_1} \cdots x_{n-1}^{\nu_{n-1}}$  such that  $\nu_1 + \cdots + \nu_{n-1} = k+1$ ,  $\nu_1, \dots, \nu_{n-1} \geq 0$ . One of our goals is to determine the annihilating ideal of the fusion product (2.2) in  $U(\mathfrak{a}[t]) = \mathbb{C}[x_a[i] \mid 1 \leq a \leq n-1, i \in \mathbb{Z}_{\geq 0}]$ . It turns out that  $x_a[i]$  belongs to this ideal if  $i$  is sufficiently large (see (2.10)), and hence we can work with polynomial rings with finitely many variables.

In order to study the inductive structure of the fusion product, it is necessary to consider a family of modules slightly more general than (2.2). In the next section, we begin by reformulating the setting to accommodate this point.

**2.2. Modules over an abelian Lie algebra.** Let  $\mathfrak{a} = \bigoplus_{a=1}^{n-1} \mathbb{C}x_a$  be an abelian Lie algebra. For  $1 \leq m \leq n$  and  $k \geq 0$ , set

$$(2.3) \quad V_k^{(m)} = \mathbb{C}[x_1, \dots, x_{n-1}] / J_k^{(m)},$$

where  $J_k^{(m)}$  is the ideal generated by the following elements:

$$(2.4) \quad J_k^{(m)} = \langle x_m, \dots, x_{n-1}, \quad x_1^{\nu_1} \cdots x_{m-1}^{\nu_{m-1}} : \nu_i \geq 0, \sum_{a=1}^{m-1} \nu_a = k+1 \rangle.$$

Notice that

$$V_k^{(m)} \simeq \bigoplus_{\nu_1 + \dots + \nu_{m-1} \leq k} \mathbb{C}x_1^{\nu_1} \cdots x_{m-1}^{\nu_{m-1}}$$

so that

$$\dim V_k^{(m)} = \begin{bmatrix} k+m-1 \\ m-1 \end{bmatrix}.$$

We regard (2.3) as a cyclic  $\mathfrak{a}$ -module with cyclic vector  $v_k^{(m)} = 1 \bmod J_k^{(m)}$ . It is a trivial module  $\mathbb{C}1$  if  $m=1$  or  $k=0$ . Note also that  $x_a = 0$  on  $V_k^{(m)}$  if  $a \geq m$ .

Let  $\mathcal{Z}$  be as in (2.1), and let

$$\mathbf{n} = (n_1, \dots, n_N), \quad \mathbf{k} = (k_1, \dots, k_N)$$

be  $N$  tuples of non-negative integers with  $1 \leq n_1, \dots, n_N \leq n$ . We consider the filtered tensor product

$$(2.5) \quad \mathcal{F}_{\mathcal{Z}}(V_{k_1}^{(n_1)}, \dots, V_{k_N}^{(n_N)}) = V_{k_1}^{(n_1)}(z_1) \otimes \cdots \otimes V_{k_N}^{(n_N)}(z_N)$$

and the fusion product

$$(2.6) \quad V_{\mathcal{Z}}(\mathbf{n}, \mathbf{k}) = V_{k_1}^{(n_1)} * \cdots * V_{k_N}^{(n_N)}(\mathcal{Z}).$$

In (2.5),  $V_k^{(n)}(z)$  denotes the evaluation module with parameter  $z$ , where  $x[i]$  acts as  $z^i x$  ( $x \in \mathfrak{a}$ ). These are cyclic  $\mathfrak{a}[t]$ -modules generated by  $v(\mathbf{n}, \mathbf{k}) = v_{k_1}^{(n_1)} \otimes \cdots \otimes v_{k_N}^{(n_N)}$ . We define a  $\mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 0}^{n-1}$  grading on (2.6) by *degree* and *weight*, with the assignment

$$\begin{aligned} \deg x_a[i] &= i, & \text{wt } x_a[i] &= (0, \dots, \overset{a\text{-th}}{1}, \dots, 0), \\ \deg v(\mathbf{n}, \mathbf{k}) &= 0, & \text{wt } v(\mathbf{n}, \mathbf{k}) &= (0, \dots, 0). \end{aligned}$$

In general, for a graded vector space  $W = \bigoplus_{d, m_1, \dots, m_{n-1}} W_{d, m_1, \dots, m_{n-1}}$ , we define its character by

$$(2.7) \quad \text{ch}_{q, z_1, \dots, z_{n-1}} W = \sum q^d z_1^{m_1} \cdots z_{n-1}^{m_{n-1}} \dim W_{d, m_1, \dots, m_{n-1}}.$$

By an inequality of characters  $\text{ch}_{q, z_1, \dots, z_{n-1}} W \leq \text{ch}_{q, z_1, \dots, z_{n-1}} W'$  we mean inequalities for all the coefficients, i.e.,  $\dim W_{d, m_1, \dots, m_{n-1}} \leq \dim W'_{d, m_1, \dots, m_{n-1}}$ .

Given  $\mathbf{n}$ , define  $X_a$  to denote the set of representations on which  $x_a$  acts nontrivially, and  $N_a$  to be the number of such representations, that is,

$$(2.8) \quad X_a = \{p \mid 1 \leq p \leq N, n_p > a\},$$

$$(2.9) \quad \mathbf{N} = (N_0, \dots, N_{n-1}), \quad N_a = |X_a|,$$

so that  $N = N_0 \geq N_1 \geq \dots \geq N_n = 0$ . We note also that if  $i \geq N_a$  then  $x_a[i] = 0$  on the fusion product (2.6). This is because

$$(2.10) \quad x_a \otimes t^{i-N_a} \prod_{p \in X_a} (t - z_p)$$

acts as 0 on the filtered tensor product (2.5).

**2.3. Annihilating ideal.** Consider the polynomial ring  $R = \mathbb{C}[S_{\mathbf{N}}]$  in indeterminates

$$(2.11) \quad S_{\mathbf{N}} = \{x_a[i] \mid 1 \leq a \leq n-1, 0 \leq i \leq N_a - 1\}.$$

We denote by  $R_{d,\mathbf{m}}$  the homogeneous component of  $R$  of degree  $d$  and weight  $\mathbf{m} = (m_1, \dots, m_{n-1})$ .

The fusion product (2.6) has a presentation as a quotient of  $R$

$$(2.12) \quad V_{\mathcal{Z}}(\mathbf{n}, \mathbf{k}) = R/I_{\mathcal{Z}}(\mathbf{n}, \mathbf{k})$$

by some homogeneous ideal  $I_{\mathcal{Z}}(\mathbf{n}, \mathbf{k})$ . In this subsection, we will find a set of elements in this ideal which are independent of  $\mathcal{Z}$  (Proposition 2.1). Later we show that they generate the whole ideal  $I_{\mathcal{Z}}(\mathbf{n}, \mathbf{k})$  (Theorem 2.4).

Set

$$(2.13) \quad x_a(z) = \sum_{i=0}^{N_a-1} x_a[i] z^{N_a-1-i}.$$

For  $\boldsymbol{\nu} = (\nu_1, \dots, \nu_{n-1}) \in \mathbb{Z}_{\geq 0}^{n-1}$ , define

$$(2.14) \quad \mu(\boldsymbol{\nu}, \mathbf{k}) = \sum_{p=1}^N \left( \sum_{a=1}^{n_p-1} \nu_a - k_p \right)_+$$

where  $u_+ = \max(u, 0)$ . Consider the ideal  $J(\mathbf{n}, \mathbf{k})$  of  $R$  generated as:

$$(2.15) \quad J(\mathbf{n}, \mathbf{k}) = \langle \text{coefficients of } z^j \text{ in } \prod_{a=1}^{n-1} x_a(z)^{\nu_a}, \forall j < \mu(\boldsymbol{\nu}, \mathbf{k}), \boldsymbol{\nu} \in \mathbb{Z}_{\geq 0}^{n-1} \rangle.$$

**Proposition 2.1.** *We have*

$$J(\mathbf{n}, \mathbf{k}) \subset I_{\mathcal{Z}}(\mathbf{n}, \mathbf{k}).$$

*That is, the following relations hold in  $R/I_{\mathcal{Z}}(\mathbf{n}, \mathbf{k})$ :*

$$(2.16) \quad \text{ord}_z \prod_{a=1}^{n-1} x_a(z)^{\nu_a} \geq \mu(\boldsymbol{\nu}, \mathbf{k}).$$

Above, for a polynomial  $\varphi(z) = \sum_{j \geq 0} c_j z^j$  we write  $\text{ord}_z \varphi \geq j_0$  if  $c_j = 0$  for  $j < j_0$ .

To prove Proposition 2.1, we prepare a few Lemmas. For each  $0 \leq a \leq n-1$ , set

$$\begin{aligned} \tilde{x}_a(z) &= \sum_{i=0}^{N_a-1} \tilde{x}_a[i] z^{N_a-1-i}, \\ \tilde{x}_a[i] &= x_a[i] - x_a[i-1]\sigma_{a,1} + x_a[i-2]\sigma_{a,2} - \cdots \\ &= \sum_{s=0}^i (-1)^s x_a[i-s]\sigma_{a,s}, \end{aligned}$$

where  $\sigma_{a,s}$  denotes the  $s$ -th elementary symmetric polynomial in the variables  $\{z_p\}_{p \in X_a}$ . Let  $x_a^{(p)}$  stand for the action of  $x_a$  on the  $p$ -th tensor component.

**Lemma 2.2.** *There exists a constant  $c_a^{(p)} \neq 0$  such that on (2.5) we have*

$$(2.17) \quad \tilde{x}_a(z_p) = c_a^{(p)} x_a^{(p)} \quad (1 \leq a \leq n_p - 1).$$

*Proof.* Let  $v$  be a vector of (2.5). Then we have

$$\begin{aligned} \sum_{i=0}^{\infty} z^{-i} \tilde{x}_a[i] v &= \prod_{p \in X_a} (1 - z_p/z) \sum_{i=0}^{\infty} z^{-i} x_a[i] v \\ &= \prod_{p \in X_a} (1 - z_p/z) \sum_{p \in X_a} \frac{1}{1 - z_p/z} x_a^{(p)} v \\ &= \sum_{p \in X_a} \prod_{\substack{p' \neq p \\ p' \in X_a}} (1 - z_{p'}/z) \cdot x_a^{(p)} v. \end{aligned}$$

In the second line, we used  $x_a^{(p)} v = 0$  for  $p \notin X_a$ . We have shown that  $\tilde{x}_a[i] v = 0$  for  $i \geq N_a$ . Multiplying  $z^{N_a-1}$  and setting  $z = z_p$ , we obtain (2.17) with  $c_a^{(p)} = \prod_{\substack{p' \neq p \\ p' \in X_a}} (z_p - z_{p'})$ .  $\square$

Let us present the filtered tensor product as a quotient of  $R$  by some ideal  $\tilde{I}_Z(\mathbf{n}, \mathbf{k})$ ,

$$\mathcal{F}_Z(V_{k_1}^{(n_1)}, \dots, V_{k_N}^{(n_N)}) = R / \tilde{I}_Z(\mathbf{n}, \mathbf{k}).$$

Recall from (2.4) that

$$\prod_{a=1}^{n_p-1} (x_a^{(p)})^{\nu_a} \in J_{k_p}^{(n_p)} \quad \text{if} \quad \sum_{a=1}^{n_p-1} \nu_a > k_p.$$

We will now interpret this condition in terms of  $x_a[i]$  by using the current  $x_a(z)$ .

Denote by  $R_d = \oplus_{\mathbf{m}} R_{d,\mathbf{m}}$  the homogeneous component of degree  $d$ , and set  $R_{\leq d} = R_0 + \cdots + R_d$ .



**Lemma 2.3.** (i) The ideal  $I_{\mathcal{Z}}(\mathbf{n}, \mathbf{k})$  consists of homogeneous elements  $P \in R_d$  ( $d \geq 0$ ), such that

$$P \equiv P' \pmod{R_{\leq d-1}} \quad \text{for some } P' \in \tilde{I}_{\mathcal{Z}}(\mathbf{n}, \mathbf{k}) \cap R_{\leq d}.$$

(ii) For each  $p$ , the elements

$$\prod_{a=1}^{n_p-1} \tilde{x}_a(z_p)^{\nu_p}, \quad (\boldsymbol{\nu} \in \mathbb{Z}_{\geq 0}^{n-1}, \sum_{a=1}^{n_p-1} \nu_a > k_p)$$

belong to the ideal  $\tilde{I}_{\mathcal{Z}}(\mathbf{n}, \mathbf{k})$ .

*Proof.* Assertion (i) is clear from the definition of the fusion product as the adjoint graded space of the filtered tensor product. Assertion (ii) is an immediate consequence of Lemma 2.2 and 2.4.  $\square$

*Proof of Proposition 2.1.* Let us prove the relation (2.16). Fix  $\boldsymbol{\nu}$ , and consider the polynomial

$$f(z) = \prod_{a=1}^{n-1} \tilde{x}_a(z)^{\nu_a} \in R[z].$$

From Lemma 2.3 (ii), we have for each  $p = 1, \dots, N$

$$(2.18) \quad \left. \frac{d^j f(z)}{dz^j} \right|_{z=z_p} \in \tilde{I}_{\mathcal{Z}}(\mathbf{n}, \mathbf{k}) \quad (0 \leq j \leq \mu_p - 1),$$

where

$$\mu_p = \left( \sum_{a=1}^{n_p-1} \nu_a - k_p \right)_+.$$

Using (2.18) and the fact that  $z_1, \dots, z_N$  are distinct, we can find polynomials  $g(z), h(z) \in R[z]$  such that

$$(2.19) \quad f(z) = \prod_{p=1}^N (z - z_p)^{\mu_p} \cdot g(z) + h(z),$$

$$h(z) \in \tilde{I}_{\mathcal{Z}}(\mathbf{n}, \mathbf{k})[z], \quad \deg h < \deg g.$$

Note that they have the form

$$f(z) = \sum_{j=0}^d a_j z^{d-j}, \quad g(z) = \sum_{j=0}^{d'} b_j z^{d'-j}, \quad h(z) = \sum_{j=d-d'+1}^d c_j z^{d-j},$$

where  $a_j, b_j, c_j \in R_{\leq j}$ .

Let  $\varphi_t$  be the automorphism of  $R[t, t^{-1}]$  defined by  $\varphi_t(b) = t^{-i}b$  for  $b \in R_i$ . Because of the above form,  $t^d \varphi_t(f)(z/t)$ ,  $t^{d'} \varphi_t(g)(z/t)$  and  $t^d \varphi_t(h)(z/t)$

have a well-defined limit as  $t \rightarrow 0$ . Denote the limit by  $f^*(z), g^*(z), h^*(z)$ , respectively. Clearly we have

$$f^*(z) = \prod_{a=1}^{n-1} x_a(z)^{\nu_a}, \quad h^*(z) \in I_{\mathcal{Z}}(\mathbf{n}, \mathbf{k}).$$

From (2.19) we obtain

$$f^*(z) = z^{\sum_{p=1}^N \mu_p} g^*(z) + h^*(z).$$

The proof is over.  $\square$

**2.4. An isomorphism.** Consider now the module

$$(2.20) \quad W(\mathbf{n}, \mathbf{k}) = W \begin{pmatrix} n_1 & \cdots & n_N \\ k_1 & \cdots & k_N \end{pmatrix} = R/J(\mathbf{n}, \mathbf{k}).$$

The following relations are immediate from the definition.

$$(2.21) \quad W \begin{pmatrix} \cdots & n_{p-1} & 1 & n_{p+1} & \cdots \\ \cdots & k_{p-1} & k_p & k_{p+1} & \cdots \end{pmatrix} = W \begin{pmatrix} \cdots & n_{p-1} & n_{p+1} & \cdots \\ \cdots & k_{p-1} & k_{p+1} & \cdots \end{pmatrix},$$

$$(2.22) \quad W \begin{pmatrix} \cdots & n_p & \cdots & n_{p'} & \cdots \\ \cdots & k_p & \cdots & k_{p'} & \cdots \end{pmatrix} = W \begin{pmatrix} \cdots & n_{p'} & \cdots & n_p & \cdots \\ \cdots & k_{p'} & \cdots & k_p & \cdots \end{pmatrix}.$$

We have also a canonical isomorphism

$$(2.23) \quad W \begin{pmatrix} \cdots & n_{p-1} & n_p & n_{p+1} & \cdots \\ \cdots & k_{p-1} & 0 & k_{p+1} & \cdots \end{pmatrix} \simeq W \begin{pmatrix} \cdots & n_{p-1} & n_{p+1} & \cdots \\ \cdots & k_{p-1} & k_{p+1} & \cdots \end{pmatrix}.$$

To see this note that if  $k_p = 0$ , then for all  $a < n_p$ , the generator  $x_a[N_a - 1]$  belongs to the ideal  $J(\mathbf{n}, \mathbf{k})$  (take  $\nu_b = \delta_{ab}$ ). Therefore, in the presentation (2.20), we can replace  $R$  by dropping all such  $x_a[N_a - 1]$ . The result is the presentation in the right hand side. In what follows we identify modules which are related by (2.23).

Our first goal is the following result.

**Theorem 2.4.** *We have an isomorphism*

$$(2.24) \quad W(\mathbf{n}, \mathbf{k}) \xrightarrow{\sim} V_{\mathcal{Z}}(\mathbf{n}, \mathbf{k})$$

*given by the canonical surjection. In particular, the fusion product is independent of the choice of  $\mathcal{Z}$ .*

We prove Theorem 2.4 in subsection 2.8.

**2.5. Subquotient modules.** In order to determine the structure of  $W(\mathbf{n}, \mathbf{k})$ , we study its subquotients. If  $k_p = 0$  for all  $p$  such that  $n_p \geq 2$ , then  $W(\mathbf{n}, \mathbf{k}) \simeq \mathbb{C}$ . Otherwise, we can find a number  $p$  satisfying the following conditions.

$$(2.25) \quad n_p \geq 2, \quad k_p > 0, \quad k_s \leq k_p \text{ for } n_s \geq n_p.$$

In what follows we fix  $p$  subject to the conditions (2.25) and set

$$(2.26) \quad a = n_p - 1, \quad 1 \leq a \leq n - 1.$$

Define

$$\begin{aligned}\mathbf{n}' &= (n_1, \dots, n_p - 1, \dots, n_N), \\ \mathbf{N}' &= (N_0, \dots, N_a - 1, \dots, N_{n-1}), \\ \mathbf{k}'' &= (k_1, \dots, k_p - 1, \dots, k_N).\end{aligned}$$

We set  $S = S_{\mathbf{N}}$ ,  $S' = S_{\mathbf{N}'}$ ,  $S'' = S_{\mathbf{N}}$ ,  $J = J(\mathbf{n}, \mathbf{k})$ ,  $J' = J(\mathbf{n}', \mathbf{k})$ ,  $J'' = J(\mathbf{n}, \mathbf{k}'')$  and

$$\begin{aligned}W' &= W \begin{pmatrix} n_1 & \cdots & n_p - 1 & \cdots & n_N \\ k_1 & \cdots & k_p & \cdots & k_N \end{pmatrix} = \mathbb{C}[S']/J', \\ W &= W \begin{pmatrix} n_1 & \cdots & n_p & \cdots & n_N \\ k_1 & \cdots & k_p & \cdots & k_N \end{pmatrix} = \mathbb{C}[S]/J, \\ W'' &= W \begin{pmatrix} n_1 & \cdots & n_p & \cdots & n_N \\ k_1 & \cdots & k_p - 1 & \cdots & k_N \end{pmatrix} = \mathbb{C}[S'']/J''.\end{aligned}$$

Our second goal is the following theorem.

**Theorem 2.5.** *Notation being as above, there exists an exact sequence*

$$(2.27) \quad 0 \longrightarrow W' \xrightarrow{\iota} W \xrightarrow{\psi} W'' \longrightarrow 0,$$

where the maps shift the grading as

$$\begin{aligned}\iota \left( W'_{d-m_a, m_1, \dots, m_a, \dots, m_{n-1}} \right) &\subset W_{d, m_1, \dots, m_a, \dots, m_{n-1}}, \\ \psi \left( W_{d, m_1, \dots, m_a, \dots, m_{n-1}} \right) &\subset W''_{d, m_1, \dots, m_a-1, \dots, m_{n-1}}.\end{aligned}$$

The maps  $\iota$  and  $\psi$  are described as follows.

Let  $\tilde{\iota}$  be the map

$$\tilde{\iota} : \mathbb{C}[S'] \rightarrow \mathbb{C}[S], \quad x_b[i] \mapsto x_b[i + \delta_{ab}].$$

In subsection 2.6 we show that  $\tilde{\iota}(J') \subset J$  (Proposition 2.6). The map  $\iota$  is the induced map

$$(2.28) \quad \iota : W' = \mathbb{C}[S']/J' \longrightarrow W = \mathbb{C}[S]/J.$$

Set  $\overline{W}' = \iota(W') \subset W$ . In other words,  $\overline{W}'$  is the subspace spanned by all monomials in  $x_b[i]$  ( $b \neq a$ ,  $0 \leq i \leq N_b - 1$ ) and  $x_a[1], \dots, x_a[N_a - 1]$ . The quotient  $W/\overline{W}'$  is spanned by monomials which are divisible by  $x_a[0]$ .

In subsection 2.7, we show that there is a well-defined map

$$(2.29) \quad \phi : W'' \longrightarrow W/\overline{W}'$$

given by

$$x_a[0]^i b \mapsto \frac{1}{i+1} x_a[0]^{i+1} b,$$

where  $b$  runs over monomials not divisible by  $x_a[0]$ . We show moreover that (2.29) is an isomorphism. The map  $\psi$  is the composition  $\phi^{-1} \circ \pi$ , where  $\pi : W \rightarrow W/\overline{W}'$  signifies the canonical surjection.

Theorem 2.5 will be proved in subsection 2.8 simultaneously with Theorem 2.4.

**2.6. The map  $\iota$ .** Here we show that the map (2.28) is well-defined.

**Proposition 2.6.** *We have  $\tilde{\iota}(J') \subset J$ .*

*Proof.* Set

$$x'_b(z) = \sum_{i=0}^{N'_b-1} x_b[i] z^{N'_b-1-i}, \quad N'_b = N_b - \delta_{ab}.$$

Then  $\tilde{\iota}(x'_b(z)) = x_b(z) - \delta_{ab}x_a[0]z^{N_a-1}$ , so that we have

$$(2.30) \quad \begin{aligned} & \tilde{\iota} \left( \prod_{b=1}^{n-1} x'_b(z)^{\nu_b} \right) \\ &= \sum_{\mu=0}^{\nu_a} (-x_a[0])^\mu \prod_{b(\neq a)} x_b(z)^{\nu_b} \cdot x_a(z)^{\nu_a-\mu} z^{(N_a-1)\mu}. \end{aligned}$$

Let  $C(j)$  denote the coefficient of  $z^j$  in  $\prod_{b=1}^{n-1} x'_b(z)^{\nu_b}$ . We are to show that  $\tilde{\iota}(C(j))$  belongs to  $J$  if  $j < d_1$ , where

$$d_1 = \sum_{s=1}^N \left( \sum_{b=1}^{n'_s-1} \nu_b - k_s \right)_+, \quad n'_s = n_s - \delta_{sp}.$$

Let  $C(\mu, j)$  denote the coefficient of  $z^j$  in the  $\mu$ -th term in the right hand side of (2.30). It belongs to  $J$  if  $j < d_2$ , where (see (2.8) and (2.9) for  $X_a$ )

$$d_2 = \sum_{s \in X_a} \left( \sum_{b=1}^{n_s-1} \nu_b - \mu - k_s \right)_+ + \sum_{s \notin X_a} \left( \sum_{b=1}^{n_s-1} \nu_b - k_s \right)_+ + (N_a - 1)\mu.$$

Noting that  $x_+ + y \geq (x + y)_+$  holds for  $y \geq 0$ ,  $a = n_p - 1$ , and that  $\mu \leq \nu_a = \nu_{n_p-1}$  implies

$$\sum_{b=1}^{n_p-1} \nu_b - \mu - k_p \geq \sum_{b=1}^{n_p-2} \nu_b - k_p = \sum_{b=1}^{n'_p-1} \nu_b - k_p,$$

we find  $d_2 \geq d_1$ . The assertion follows from this.  $\square$

**2.7. Dual space.** In order to construct the map  $\phi$  in (2.29), we pass to the dual space  $W(\mathbf{n}, \mathbf{k})^*$  and use a realization for the latter in terms of polynomials.

Recall  $R = \mathbb{C}[S_{\mathbf{N}}]$ , with  $S_{\mathbf{N}} = \{x_b[i] \mid 0 \leq i \leq N_b - 1, 1 \leq b \leq n - 1\}$ . Consider the restricted dual space

$$R^* = \bigoplus_{d, \mathbf{m}} (R_{d, \mathbf{m}})^*.$$

Each homogeneous component  $(R_{d,\mathbf{m}})^*$  is realized as a subspace of the ring of polynomials  $\mathbb{C}[T_{\mathbf{m}}]$  in indeterminates

$$T_{\mathbf{m}} = \{y_{bj} \mid 1 \leq j \leq m_b, 1 \leq b \leq n-1\},$$

where  $\mathbf{m} = (m_1, \dots, m_{n-1})$ . In this space we count the total degree  $d^*$  by assigning  $\deg y_{bj} = 1$ . Let  $\mathcal{F}_{d^*, \mathbf{m}, \mathbf{N}}$  denote the subspace of  $\mathbb{C}[T_{\mathbf{m}}]$  consisting of all polynomials  $f$  with the properties

- (i)  $f$  is symmetric in each group of variables  $(y_{b1}, \dots, y_{bm_b})$ ,
- (ii)  $f$  has total degree  $d^*$ , and  $\deg_{y_{bj}} f < N_b$  for each  $b, j$ .

Set

$$d(\mathbf{N}, \mathbf{m}) = \sum_{b=1}^{n-1} (N_b - 1)m_b.$$

We have an isomorphism

$$(2.31) \quad R_{d,m}^* \xrightarrow{\sim} \mathcal{F}_{d^*, \mathbf{m}, \mathbf{N}}, \quad d + d^* = d(\mathbf{N}, \mathbf{m})$$

given by

$$\bigoplus_{d=0}^{d(\mathbf{N}, \mathbf{m})} \mathbb{C}[S_{\mathbf{N}}]_{d, \mathbf{m}}^* \ni \theta \mapsto \theta \left( \prod_{\substack{1 \leq b \leq n-1 \\ 1 \leq j \leq m_b}} x_b(y_{bj}) \right) \in \bigoplus_{d^*=0}^{d(\mathbf{N}, \mathbf{m})} \mathcal{F}_{d^*, \mathbf{m}, \mathbf{N}}.$$

With respect to this pairing, monomials

$$\prod_{a=1}^{n-1} x_a[i_1^{(a)}] \cdots x_a[i_{m_a}^{(a)}] \quad (0 \leq i_1^{(a)} \leq \cdots \leq i_{m_a}^{(a)} \leq N_a - 1)$$

and

$$\prod_{a=1}^{n-1} \text{Sym}(y_{a1}^{N_a-1-i_1^{(a)}} \cdots y_{am_a}^{N_a-1-i_{m_a}^{(a)}})$$

are dual bases to each other, where the symbol  $\text{Sym}$  stands for the normalized sum over all permutations of  $i_1^{(a)}, \dots, i_{m_a}^{(a)}$ .

We have introduced the quotient space  $W(\mathbf{n}, \mathbf{k})$  of  $R = \mathbb{C}[S_{\mathbf{N}}]$  by the homogeneous ideal  $J = J(\mathbf{n}, \mathbf{k})$  (2.15). The dual space  $(W(\mathbf{n}, \mathbf{k})_{d, \mathbf{m}})^*$  is the orthogonal complement to  $J$  under the coupling given above. Explicitly it is described as follows. Given  $\boldsymbol{\nu}$  such that  $\nu_b \leq m_b$  ( $1 \leq b \leq n-1$ ), we consider the specialization of variables

$$(2.32) \quad \begin{aligned} y_{11} &= \cdots = y_{1\nu_1} = z \\ &\vdots \\ y_{n-1,1} &= \cdots = y_{n-1,\nu_{n-1}} = z \end{aligned}$$

and denote by  $\Phi(f, \nu_1, \dots, \nu_{n-1}; z)$  the polynomial obtained from  $f$  by (2.32).

**Lemma 2.7.** *The dual space  $(W(\mathbf{n}, \mathbf{k})_{d, \mathbf{m}})^*$  is isomorphic to the subspace of  $\mathcal{F}_{d^*, \mathbf{m}, \mathbf{N}}$  consisting of polynomials  $f$  satisfying (see (2.14) for  $\mu(\boldsymbol{\nu}, \mathbf{k})$ )*

$$(2.33) \quad \text{ord}_z \Phi(f, \nu_1, \dots, \nu_{n-1}; z) \geq \mu(\boldsymbol{\nu}, \mathbf{k})$$

for all  $\boldsymbol{\nu} \in \mathbb{Z}_{\geq 0}^{n-1}$  such that  $\nu_b \leq m_b$  ( $1 \leq b \leq n-1$ ). Namely,  $f \in \mathcal{F}_{d^*, \mathbf{m}, \mathbf{N}}$  is orthogonal to  $J$  if and only if it satisfies the condition (2.33).

We need also the orthogonal complement to the subspace  $\text{Im } \tilde{t}$ . An element  $f \in \mathcal{F}_{d^*, \mathbf{m}, \mathbf{N}}$  is orthogonal to  $\text{Im } \tilde{t}$  if and only if

$$(2.34) \quad \text{it does not contain monomials of the form } \prod_{\substack{1 \leq b \leq n-1 \\ 1 \leq j \leq m_b}} y_{bj}^{i_{bj}}$$

$$\text{with } i_{a1}, \dots, i_{am_a} < N_a - 1.$$

Such an  $f$  can be written uniquely in the form

$$(2.35) \quad f = \sum_{j=1}^{m_a} y_{aj}^{N_a-1} g^{(j)}$$

with some polynomial  $g \in \mathbb{C}[T_{m_1, \dots, m_a-1, \dots, m_{n-1}}]$ , where

$$g^{(j)} = g(y_{11}, \dots; \dots, \widehat{y_{aj}}, \dots; \dots, y_{n-1, m_{n-1}}).$$

In the right hand side the variable  $y_{aj}$  is omitted. Let  $\mathcal{F}'_{d^*, \mathbf{m}, \mathbf{N}}$  be the subspace of polynomials satisfying (2.34). Setting  $\tilde{\phi}^*(f) = g$ , we obtain an injective map

$$\tilde{\phi}^* : \mathcal{F}'_{d^*, m_1, \dots, m_a, \dots, m_{n-1}, \mathbf{N}} \longrightarrow \mathcal{F}_{d^*-N_a+1, m_1, \dots, m_a-1, \dots, m_{n-1}, \mathbf{N}}.$$

Note that the dual space  $(W/\overline{W}')^*_{d^*, \mathbf{m}}$  is isomorphic to the subspace of  $\mathcal{F}'_{d^*, \mathbf{m}, \mathbf{N}}$  which is orthogonal to  $J$ .

**Proposition 2.8.** *If  $f \in \mathcal{F}'_{d^*, \mathbf{m}, \mathbf{N}}$  is orthogonal to  $J$ , then  $g = \tilde{\phi}^*(f)$  is orthogonal to  $J''$ . Hence  $\tilde{\phi}^*$  induces an injection*

$$\phi^* : (W/\overline{W}')^*_{d^*, m_1, \dots, m_a, \dots, m_{n-1}} \longrightarrow (W'')^*_{d^*-N_a+1, m_1, \dots, m_a-1, \dots, m_{n-1}}.$$

*Proof.* By Lemma 2.7,  $f$  satisfies the condition (2.33). We are to show that

$$(2.36) \quad \text{ord}_z \Phi(g, \nu_1, \dots, \nu_{n-1}; z) \geq \sum_{s=1}^N \left( \sum_{1 \leq b \leq n_s-1} \nu_b - k_s + \delta_{sp} \right)_+$$

holds for any  $\boldsymbol{\nu}$  with  $\nu_b \leq m_b - \delta_{ab}$  ( $1 \leq b \leq n-1$ ).

Specializing (2.35), we obtain

$$(2.37) \quad \begin{aligned} & \Phi(f, \nu_1, \dots, \nu_a, \dots, \nu_{n-1}; z) \\ &= \nu_a z^{N_a-1} \Phi(g, \nu_1, \dots, \nu_a - 1, \dots, \nu_{n-1}; z) \\ &+ \sum_{\nu_a+1 \leq j \leq m_a} y_{aj}^{N_a-1} \Phi(g^{(j)}, \nu_1, \dots, \nu_a, \dots, \nu_{n-1}; z). \end{aligned}$$

We distinguish two cases.

**Case**  $\sum_{1 \leq b \leq a} \nu_b \leq k_p - 1$ : In this case, the term with  $s = p$  in the right hand side of (2.36) does not contribute. If  $\nu_a = 0$ , then (2.36) follows from (2.33), (2.37) and the uniqueness of the representation (2.35). The general case follows by ascending induction on  $\nu_a$ . Namely, we apply the induction hypothesis (2.36) for the first term in the right hand side of (2.37), and conclude (2.36) for the second term.

**Case**  $\sum_{1 \leq b \leq a} \nu_b \geq k_p$ : We use (2.37) by changing  $\nu_a$  to  $\nu_a + 1$ :

$$\begin{aligned} & \Phi(f, \nu_1, \dots, \nu_a + 1, \dots, \nu_{n-1}; z) \\ &= (\nu_a + 1) z^{N_a - 1} \Phi(g, \nu_1, \dots, \nu_a, \dots, \nu_{n-1}; z) \\ &+ \sum_{\nu_a + 2 \leq j \leq m_a} y_{aj}^{N_a - 1} \Phi(g^{(j)}, \nu_1, \dots, \nu_a + 1, \dots, \nu_{n-1}; z). \end{aligned}$$

Consider first the case  $\nu_a = m_a - 1$ . The second term in the right hand side is then void. We have

$$\begin{aligned} & \text{ord}_z \Phi(g, \nu_1, \dots, \nu_a, \dots, \nu_{n-1}; z) \\ (2.38) \geq & \sum_{s \notin X_a} \left( \sum_{1 \leq b \leq n_s - 1} \nu_b - k_s \right)_+ + \sum_{s \in X_a} \left( \sum_{1 \leq b \leq n_s - 1} \nu_b + 1 - k_s \right)_+ - (N_a - 1). \end{aligned}$$

Since  $a = n_p - 1$ ,  $s \in X_a$  implies  $n_p \leq n_s$ , and hence

$$\sum_{1 \leq b \leq n_s - 1} \nu_b \geq \sum_{1 \leq b \leq a} \nu_b \geq k_p \geq k_s.$$

In the last step we used the assumption (2.25). Therefore we find that the right hand side of (2.38) is not less than

$$\sum_{s=1}^N \left( \sum_{1 \leq b \leq n_s - 1} \nu_b - k_s + \delta_{sp} \right)_+.$$

The general case follows by descending induction on  $\nu_a$ . Namely, we apply the induction hypothesis (2.36) for the second term in the right hand side of (2.37), and conclude (2.36) for the first term.  $\square$

Let

$$\tilde{\phi} : \mathbf{C}[S_{\mathbf{N}}]_{d, m_1, \dots, m_a - 1, \dots, m_{n-1}} \longrightarrow (\mathbf{C}[S_{\mathbf{N}}]/\text{Im } \tilde{\iota})_{d, \mathbf{m}}$$

be the map dual to  $\tilde{\phi}^*$ . Decomposing  $\mathbf{C}[S_{\mathbf{N}}]_{d, m_1, \dots, m_a - 1, \dots, m_{n-1}}$  as

$$\oplus_{i=0}^{m_a - 1} x_a[0]^i (\text{Im } \tilde{\iota})_{d, m_1, \dots, m_a - 1 - i, \dots, m_{n-1}},$$

we can explicitly write  $\tilde{\phi}$  as

$$\tilde{\phi} : x_a[0]^i (\text{Im } \iota)_{d, m_1, \dots, m_a - 1 - i, \dots, m_{n-1}} \ni b \mapsto \frac{x_a[0]}{i + 1} b \in (\mathbf{C}[S_{\mathbf{N}}]/\text{Im } \tilde{\iota})_{d, \mathbf{m}}.$$

We have shown in Proposition 2.8 that if  $b \in \mathbf{C}[S_{\mathbf{N}}]_{d, m_1, \dots, m_a - 1, \dots, m_{n-1}}$  belongs to  $J''$  then the image  $\tilde{\phi}(b)$  belongs to  $J$ . Therefore,  $\tilde{\phi}$  induces the surjection  $\phi$  of (2.29).

**2.8. Proof of Theorems 2.4, 2.5.** Summarizing the results of subsections 2.6 and 2.7, we obtain the diagram

$$\begin{array}{ccccccc} & & W' & & & & W'' \\ & & \downarrow \iota & & & & \downarrow \phi \\ 0 & \longrightarrow & \overline{W}' & \longrightarrow & W & \xrightarrow{\pi} & W/\overline{W}' \longrightarrow 0 \end{array}$$

where the vertical arrows are surjective. In particular we have

$$(2.39) \quad \dim W \leq \dim W' + \dim W''.$$

We are now in a position to finish the proof of Theorems 2.4, 2.5. Set

$$\begin{aligned} \tilde{d}(\mathbf{n}, \mathbf{k}) &= \dim W(\mathbf{n}, \mathbf{k}), \\ d(\mathbf{n}, \mathbf{k}) &= \dim V_{\mathbb{Z}}(\mathbf{n}, \mathbf{k}). \end{aligned}$$

From (2.24) we have  $\tilde{d}(\mathbf{n}, \mathbf{k}) \geq d(\mathbf{n}, \mathbf{k})$ .

On the other hand, under the assumption (2.25), we have from (2.39) the inequality

$$(2.40) \quad \tilde{d} \begin{pmatrix} \cdots & a+1 & \cdots \\ \cdots & k_p & \cdots \end{pmatrix} \leq \tilde{d} \begin{pmatrix} \cdots & a & \cdots \\ \cdots & k_p & \cdots \end{pmatrix} + \tilde{d} \begin{pmatrix} \cdots & a+1 & \cdots \\ \cdots & k_p-1 & \cdots \end{pmatrix}.$$

We have also the relations

$$\begin{aligned} \tilde{d} \begin{pmatrix} 1, & \cdots, & 1 \\ k_1, & \cdots, & k_N \end{pmatrix} &= 1, \\ \tilde{d} \begin{pmatrix} \cdots & n_p & \cdots \\ \cdots & 0 & \cdots \end{pmatrix} &= \tilde{d} \begin{pmatrix} \cdots & n_p-1 & \cdots \\ \cdots & 0 & \cdots \end{pmatrix}, \\ \tilde{d} \begin{pmatrix} \cdots & n_p & \cdots & n_{p'} & \cdots \\ \cdots & k_p & \cdots & k_{p'} & \cdots \end{pmatrix} &= \tilde{d} \begin{pmatrix} \cdots & n_{p'} & \cdots & n_p & \cdots \\ \cdots & k_{p'} & \cdots & k_p & \cdots \end{pmatrix}. \end{aligned}$$

The above relations are also true for  $d(\mathbf{n}, \mathbf{k})$ , and moreover the equality holds in (2.40), due to the identity for binomial coefficients. These equations determine  $d(\mathbf{n}, \mathbf{k})$  uniquely. Therefore by induction we have for all  $\mathbf{n}, \mathbf{k}$  that

$$\tilde{d}(\mathbf{n}, \mathbf{k}) \leq d(\mathbf{n}, \mathbf{k}).$$

This implies that in (2.40) the equality always takes place. Hence  $\iota$  and  $\phi$  are isomorphisms.  $\square$

**2.9. Characters and  $q$ -supernomial coefficients.** In this section we identify the characters of the fusion product with the  $q$ -supernomial coefficients for  $\mathfrak{sl}_n$  known in the literature [HKKOTY], [SW2]. Our consideration is restricted to the case of symmetric tensor representations.

For the description of the characters, we find it convenient to use the language of partitions. For a partition  $\lambda = (\lambda_1, \dots, \lambda_N)$ , let  $\lambda'$  denote the conjugate partition. We write  $|\lambda| = \sum_{i=1}^N \lambda_i$ . We identify partitions



obtained by concatenating 0's at the end. Reordering  $n_p, k_p$  by using (2.22), we can assume without loss of generality that

$$(2.41) \quad n_1 \geq \cdots \geq n_N,$$

$$(2.42) \quad k_{N_a+1} \geq \cdots \geq k_{N_{a-1}} \quad (a = 1, \dots, n).$$

Define partitions  $\mu^{(1)} \subset \cdots \subset \mu^{(n)}$  by setting

$$\begin{aligned} \mu^{(a)} &= \sum_{b=1}^a \kappa^{(b)}, \\ \kappa^{(b)} &= (k_{N_b+1}, \dots, k_{N_{b-1}})'. \end{aligned}$$

Note that  $\mu_1^{(a)} = N - N_a$ . We adopt the following notation

$$(2.43) \quad W^{(n)}[\mu^{(n)}, \dots, \mu^{(1)}] = W(\mathbf{n}, \mathbf{k}).$$

When  $k_1 = \cdots = k_{N_{n-1}} = 0$ , we have  $\mu^{(n)} = \mu^{(n-1)}$ . In this case we understand that

$$W^{(n)}[\mu^{(n-1)}, \mu^{(n-1)}, \dots, \mu^{(1)}] = W^{(n-1)}[\mu^{(n-1)}, \dots, \mu^{(1)}].$$

Assuming  $\mu^{(n)} \neq \mu^{(n-1)}$  and therefore  $n_1 = n$ , let  $k = k_1$ . Theorem 2.5 implies the following exact sequence.

$$\begin{aligned} (2.44) \quad 0 &\longrightarrow W^{(n)}[\mu^{(n)}, \mu^{(n-1)} + (1^k), \dots, \mu^{(1)}] \\ &\xrightarrow{\iota} W^{(n)}[\mu^{(n)}, \mu^{(n-1)}, \dots, \mu^{(1)}] \xrightarrow{\psi} W^{(n)}[\mu^{(n)} - \mathbf{e}_k, \mu^{(n-1)}, \dots, \mu^{(1)}] \longrightarrow 0. \end{aligned}$$

Here  $\mathbf{e}_k$  denotes the unit vector  $(\delta_{jk})_{1 \leq j \leq k}$  and  $(1^k) = \sum_{i=1}^k \mathbf{e}_i$ . The subspace corresponds to changing  $n_1$  to  $n_1 - 1$ , while the quotient space corresponds to changing  $k_1$  to  $k_1 - 1$ .

Let us rewrite (2.44) in terms of the character (see (2.7))

$$\begin{aligned} (2.45) \quad \chi^{(n)}[\mu^{(n)}, \dots, \mu^{(1)}](q, z_1, \dots, z_{n-1}) \\ &= \text{ch}_{q, z_1, \dots, z_{n-1}} W^{(n)}[\mu^{(n)}, \dots, \mu^{(1)}] \\ &= \sum q^d z_1^{m_1} \cdots z_{n-1}^{m_{n-1}} \dim W^{(n)}[\mu^{(n)}, \dots, \mu^{(1)}]_{d, m_1, \dots, m_{n-1}}. \end{aligned}$$

**Proposition 2.9.** *We have*

$$\begin{aligned} (2.46) \quad \chi^{(n)}[\mu^{(n)}, \dots, \mu^{(1)}](q, z_1, \dots, z_{n-1}) \\ &= \chi^{(n)}[\mu^{(n)}, \mu^{(n-1)} + (1^k), \dots, \mu^{(1)}](q, z_1, \dots, qz_{n-1}) \\ &\quad + z_{n-1} \chi^{(n)}[\mu^{(n)} - \mathbf{e}_k, \dots, \mu^{(1)}](q, z_1, \dots, z_{n-1}). \end{aligned}$$

It is useful to modify the character

$$\begin{aligned} \tilde{\chi}^{(n)}[\mu^{(n)}, \dots, \mu^{(1)}](q, z_1, \dots, z_{n-1}) \\ &= \chi^{(n)}[\mu^{(n)}, \dots, \mu^{(1)}](q, q^{\mu_1^{(1)}} z_1, \dots, q^{\mu_1^{(n-1)}} z_{n-1}), \end{aligned}$$

so as to absorb the shift of variable  $z_{n-1} \rightarrow qz_{n-1}$ . Eq. (2.46) then becomes

$$(2.47) \quad \begin{aligned} & \tilde{\chi}^{(n)}[\mu^{(n)}, \dots, \mu^{(1)}] \\ &= \tilde{\chi}^{(n)}[\mu^{(n)}, \mu^{(n-1)} + (1^k), \dots, \mu^{(1)}] + q^{\mu_1^{(n-1)}} z_{n-1} \tilde{\chi}^{(n)}[\mu^{(n)} - \mathbf{e}_k, \mu^{(n-1)}, \dots, \mu^{(1)}] \end{aligned}$$

where we have suppressed the variables  $(q, z_1, \dots, z_{n-1})$  which are common to all terms. The base of the recursion is

$$\tilde{\chi}^{(1)}[\mu^{(1)}] = 1.$$

Using (2.44) repeatedly, we are led to

**Proposition 2.10.** *There exists a filtration by submodules*

$$W^{(n)}[\mu^{(n)}, \dots, \mu^{(1)}] = F^0 \supset F^1 \supset F^2 \supset \dots$$

such that each composition factor  $F^j/F^{j-1}$  has the form

$$W^{(n-1)}[\nu^{(n-1)}, \mu^{(n-2)}, \dots, \mu^{(1)}],$$

with appropriate shift of gradings, where  $\nu^{(n-1)}$  is a partition satisfying  $\mu^{(n)} \supset \nu^{(n-1)} \supset \mu^{(n-1)}$ .

In terms of the characters, it means a relation of the form

$$(2.48) \quad \begin{aligned} & \tilde{\chi}^{(n)}[\mu^{(n)}, \mu^{(n-1)}, \dots, \mu^{(1)}] \\ &= \sum_{\mu^{(n)} \supset \nu^{(n-1)} \supset \mu^{(n-1)}} C_{\nu^{(n-1)}}(q, z_{n-1}) \tilde{\chi}^{(n-1)}[\nu^{(n-1)}, \mu^{(n-2)}, \dots, \mu^{(1)}], \end{aligned}$$

where  $C_{\nu^{(n-1)}}(q, z_{n-1})$  are some polynomials in  $q$  and  $z_{n-1}$ .

*Example.* The following diagram illustrates the process of the reduction on the example  $n = 2$ ,  $k = 2$ ,  $\mu^{(2)} = (3, 2)$ ,  $\mu^{(1)} = \emptyset$  and  $\nu^{(1)} = (2, 1)$ .

$$\begin{array}{ccccc} (\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}, \emptyset) & \xrightarrow{z_1} & (\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \emptyset) \\ \uparrow & & \uparrow \\ (\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \end{array}) & \xrightarrow{qz_1} & (\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \end{array}) & \xrightarrow{qz_1} & (\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \end{array}) \\ & & \uparrow & & \uparrow \\ & & (\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}) & \xrightarrow{q^2 z_1} & (\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}) \end{array}$$

The vertical and the horizontal arrows correspond to the maps  $\iota$  and  $\psi$ , respectively. The coefficients appearing in (2.47) are as indicated.  $\square$

Let us compute the coefficients  $C_{\nu^{(n-1)}}(q, z_{n-1})$ . Assuming  $\mu^{(n)} \neq \mu^{(n-1)}$ , we apply (2.48) repeatedly for  $k_1 = k, k-1, \dots$ , and collect the terms which have the array of partitions of the form  $(\nu^{(n-1)}, \nu^{(n-1)}, \mu^{(n-2)}, \dots)$ . First consider  $k_1 = k$ . If we take the first term from (2.48)  $m_1$  times and the second term  $m_2$  times, then the resulting array of partitions becomes

$(\mu^{(n)} - m_2 \mathbf{e}_k, \mu^{(n-1)} + m_1(1^k), \mu^{(n-2)}, \dots)$ . Comparing the  $k$ -th row, we have  $m_1 = \nu_k^{(n-1)} - \mu_k^{(n-1)}$  and  $m_2 = \mu_k^{(n)} - \nu_k^{(n-1)}$ . There are altogether  $\binom{m_1 + m_2}{m_1}$  such terms. It is easy to see that the sum of the corresponding coefficients is

$$(q^{\mu_1^{(n-1)}} z)^{\mu_k^{(n)} - \nu_k^{(n-1)}} \begin{bmatrix} \mu_k^{(n)} - \mu_k^{(n-1)} \\ \nu_k^{(n-1)} - \mu_k^{(n-1)} \end{bmatrix}.$$

One can continue a similar process for  $k_1 = k - 1, \dots, 1$ . The coefficient  $C_{\nu^{(n-1)}}(q, z_{n-1})$  is the product of all factors resulting from each step.

Let us summarize the above calculation.

For a pair of partitions  $\kappa \supset \nu$ , define

$$(2.49) \quad F_{\kappa, \nu}(q) = q^{\sum_{a=1}^{k-1} \nu_{a+1}(\kappa_a - \nu_a)} \prod_{a=1}^k \begin{bmatrix} \kappa_a - \nu_{a+1} \\ \nu_a - \nu_{a+1} \end{bmatrix}.$$

Here  $k = \kappa'_1$  is the length of  $\kappa$ , and the  $q$ -binomial symbol is

$$\begin{bmatrix} m \\ n \end{bmatrix} = \begin{cases} \frac{[m]!}{[n]![m-n]!} & (0 \leq n \leq m), \\ 0 & \text{otherwise,} \end{cases}$$

with  $[n]! = \prod_{j=1}^n (1 - q^j)/(1 - q)$ . Returning to the definition (2.45), we find the following.

**Theorem 2.11.** *We have*

$$(2.50) \quad \begin{aligned} & \chi^{(n)}[\mu^{(n)}, \mu^{(n-1)}, \dots, \mu^{(1)}](q, z_1, \dots, z_{n-1}) \\ &= \sum_{\mu^{(n)} \supset \nu \supset \mu^{(n-1)}} z_{n-1}^{|\mu^{(n)}| - |\nu|} F_{\mu^{(n)} - \mu^{(n-1)}, \nu - \mu^{(n-1)}}(q) \\ & \quad \times \chi^{(n-1)}[\nu, \dots, \mu^{(1)}](q, z_1, \dots, z_{n-2}). \end{aligned}$$

Hence we obtain a fermionic expression for the character

$$(2.51) \quad \begin{aligned} & \chi^{(n)}[\mu^{(n)}, \dots, \mu^{(1)}](q, z_1, \dots, z_{n-1}) \\ &= \sum \prod_{a=1}^{n-1} \left( z_a^{|\nu^{(a+1)}| - |\nu^{(a)}|} F_{\nu^{(a+1)} - \mu^{(a)}, \nu^{(a)} - \mu^{(a)}}(q) \right), \end{aligned}$$

where the sum extends over sequences of partitions

$$\mu^{(n)} = \nu^{(n)} \supset \nu^{(n-1)} \supset \dots \supset \nu^{(1)}$$

satisfying  $\nu^{(a)} \supset \mu^{(a)}$  ( $1 \leq a \leq n-1$ ).

In the case  $n_1 = \dots = n_N = n$ , we have  $\mu^{(1)} = \dots = \mu^{(n-1)} = \emptyset$  and  $\mu^{(n)} = \mu$  where  $\mu = (k_1, \dots, k_N)'$ . By comparing (2.51) with known results in the literature, we conclude that the coefficients of monomials in  $z_a$  are the  $q$ -supernomial coefficients. More precisely, setting

$$W^{(n)}[\mu] = W^{(n)}[\mu, \emptyset, \dots, \emptyset], \quad \chi^{(n)}[\mu] = \chi^{(n)}[\mu, \emptyset, \dots, \emptyset],$$

we have

$$(2.52) \quad \chi^{(n)}[\mu](q, z_1, \dots, z_{n-1}) = \sum_{\substack{\lambda \in \mathbb{Z}_{\geq 0}^n \\ |\lambda| = |\mu|}} z_1^{\lambda_1} \cdots z_{n-1}^{\lambda_{n-1}} \tilde{S}_{\lambda, \mu}(q).$$

Here  $\lambda = (\lambda_1, \dots, \lambda_n)$  and  $\tilde{S}_{\lambda, \mu}(q)$  denotes the  $q$ -supernomial coefficient as given in formula (2.1) of [S].

The exact sequence (2.44) gives also a recursive method of constructing a monomial basis of  $W^{(n)}[\mu^{(n)}, \dots, \mu^{(1)}]$ .


**Theorem 2.12.** *There exists a basis  $B^{(n)}[\mu^{(n)}, \dots, \mu^{(1)}]$  of  $W^{(n)}[\mu^{(n)}, \dots, \mu^{(1)}]$  defined recursively as follows.*

- (i) *Each element in the set  $B^{(n)}[\mu^{(n)}, \dots, \mu^{(1)}]$  is represented by a monomial in the variables  $\{x_b[i]\}$ .*
- (ii)  $B^{(1)}[\mu^{(1)}] = \{1\}$ .
- (iii) *If the length of  $\mu^{(n)} - \mu^{(n-1)}$  is  $k > 0$ , then*

$$B^{(n)}[\mu^{(n)}, \dots, \mu^{(1)}] = \iota \left( B^{(n)}[\mu^{(n)}, \mu^{(n-1)} + (1^k), \dots, \mu^{(1)}] \right) \bigsqcup x_a[0] B^{(n)}[\mu^{(n)} - \mathbf{e}_k, \dots, \mu^{(1)}],$$

where the second term in the right hand side means elementwise multiplication by  $x_a[0]$ .

*Example.* Consider the case of  $\mathfrak{sl}_2$  (with  $e \stackrel{\text{def}}{=} x_1$ ). Computing the basis for the example above, with  $\mu^{(2)} = (3, 2)$  and  $\mu^{(1)} = \emptyset$ , we have



Here, the dotted arrows correspond to the map  $\iota : e[i] \rightarrow e[i+1]$ , and the solid arrows to multiplication by  $e[0]$ . As a result, and noting that

$B^{(2)}[\mu + (1), \mu] = \{1, e[0]\}$ , we have

$$\begin{aligned} B^{(2)}[(3, 2), \emptyset] = & \{1, e[0], e[0]^2, e[0]^3, e[0]^4, e[0]^5, e[1], e[0]e[1], \\ & e[0]^2e[1], e[0]^3e[1], e[2], e[1]^2, e[0]e[2], \\ & e[0]e[1]^2, e[0]^2e[2], e[0]^2e[1]^2, e[1]e[2], e[1]^3\}. \end{aligned}$$

### 3. CHARACTER OF $\widehat{\mathfrak{sl}}_2$ SPACES OF COINVARIANTS

In [FJKLM], we have introduced a large class of spaces of coinvariants, and established their equivalence to certain quotients of filtered tensor products of finite-dimensional modules. In this section, we restrict to the simplest case where the underlying simple Lie algebra is  $\mathfrak{g} = \mathfrak{sl}_2$ , and apply the results of the previous section to the character of the space of coinvariants.

**3.1. Spaces of coinvariants.** We retain the following notation of [FJKLM]:  $e, f, h$  are the standard generators of  $\mathfrak{sl}_2$ .  $V_l$  is the  $(l+1)$ -dimensional irreducible representation with lowest weight vector  $v_l$ .  $L_l^{(k)}(\infty)$  is the level  $k$  integrable module of the affine Lie algebra  $\widehat{\mathfrak{sl}}_2$  ‘placed at infinity’. It has  $V_l$  as its subspace of degree 0 and  $x[i]V_l = 0$  if  $x \in \mathfrak{sl}_2, i < 0$ . By  $\mathcal{V}^{(k)} = \bigoplus_{l=0}^k \mathbb{Z} \cdot [l]$  we denote the level  $k$  Verlinde algebra for  $\mathfrak{sl}_2$  with basis  $[l]$  ( $l = 0, \dots, k$ ). We write an element  $a \in \mathcal{V}^{(k)}$  as  $a = \sum_{l=0}^k (a : [l])_k [l]$ , where  $(a : [l])_k \in \mathbb{Z}$ . For an  $\mathfrak{sl}_2[t]$ -module  $L$  and a right ideal  $Y \subset U(\mathfrak{sl}_2[t])$ , we write  $L/Y$  for the space of coinvariants  $L/YL$ . Let  $\mathbf{m} = (m_1, \dots, m_k)$  be an array of non-negative integers. The level-restricted Kostka polynomial for  $\mathfrak{sl}_2$  is

$$(3.1) \quad K_{l, \mathbf{m}}^{(k)}(q) = \sum_{\substack{\mathbf{s} \in \mathbb{Z}_{\geq 0}^k \\ 2|\mathbf{s}| = |\mathbf{m}| - l}} q^{\mathbf{s} \cdot A\mathbf{s} + \mathbf{v} \cdot \mathbf{s}} \begin{bmatrix} A(\mathbf{m} - 2\mathbf{s}) - \mathbf{v} + \mathbf{s} \\ \mathbf{s} \end{bmatrix},$$

where  $A$  is a  $k \times k$  matrix with entries  $A_{ij} = \min(i, j)$ ,  $v_i = (i - k + l)_+$ ,  $|\mathbf{u}| = \sum_{i=1}^k i u_i$ , and  $\mathbf{u} \cdot \mathbf{u}' = \sum_{i=1}^k u_i u'_i$ . In what follows we use also the partition

$$(3.2) \quad \mu = (k^{m_k} \dots 1^{m_1})'$$

to label  $\mathbf{m}$ . We abuse the notation and write  $K_{l, \mu}^{(k)}(q)$  for  $K_{l, \mathbf{m}}^{(k)}(q)$ . We recall also the following formula in the Verlinde algebra  $\mathcal{V}^{(k)}$ , which will be used later.

$$(3.3) \quad [k]^{m_k} \dots [1]^{m_1} = \sum_{l=0}^k K_{l, \mu}^{(k)}(1) \cdot [l].$$

When we deal with fusion product, we consider also  $\mathfrak{sl}_2$  embedded in  $\mathfrak{sl}_3$ . We write the generators as  $e_{23}, e_{32}, h_{23}$ . In particular, we use  $h_{23}[0]$  for the action on the fusion product, and  $h[0]$  for that on  $L_l^{(k)}(\infty)$ .

Consider the fusion product of irreducible modules

$$W^{(2)}[\mu] = \overbrace{V_k * \dots * V_k}^{m_k} * \dots * \overbrace{V_1 * \dots * V_1}^{m_1}$$

taking the tensor product of lowest weight vectors

$$\overbrace{v_k \otimes \cdots \otimes v_k}^{m_k} \otimes \cdots \otimes \overbrace{v_1 \otimes \cdots \otimes v_1}^{m_1}$$

as cyclic vector. The eigenvalue of  $h_{23}[0]$  on it is  $-|\mu|$ . Denote by  $I_l^{(k)}$  the left ideal of  $U(\mathfrak{sl}_2[t])$  generated by the elements

$$h_{23}[0] + l, \quad e_{23}[0], \quad e_{23}[1]^{k-l+1}.$$

Let  $S(I_l^{(k)})$  be the right ideal obtained by applying the antipode  $S$ . The following result was obtained in [FJKLM], Theorem 4.1.

**Proposition 3.1.** *We have*

$$\text{ch}_{q,z_1} W^{(2)}[\mu]/S(I_l^{(k)}) = z_1^{\frac{|\mu|-l}{2}} K_{l,\mu}^{(k)}(q).$$

**3.2. Fusion product of  $\mathfrak{sl}_3$ -modules.** In this section, we consider reducible  $\mathfrak{sl}_2$ -modules of the form

$$(3.4) \quad W_r = V_0 \oplus \cdots \oplus V_r.$$

It is generated over  $\mathfrak{sl}_2$  by the sum of lowest weight vectors  $w_r = v_0 + \cdots + v_r$ . We make use of the fact that (3.4) can also be viewed as the  $r$ -th symmetric tensor  $S^r(V_{(1,0)})$  of the defining representation of  $\mathfrak{sl}_3$ , restricted to the subalgebra  $\mathfrak{sl}_2 \subset \mathfrak{sl}_3$ .

Let  $\mathbf{M} = (M_1, \dots, M_k) \in \mathbb{Z}_{\geq 0}^k$ , and

$$(3.5) \quad \lambda = (k^{M_k}, \dots, 1^{M_1})'.$$

Fix a set of distinct complex numbers  $\mathcal{Z} = (z_1, \dots, z_N)$ ,  $N = M_1 + \cdots + M_k$ . Consider the filtered tensor product of  $\mathfrak{sl}_2[t]$ -modules

$$(3.6) \quad \mathcal{F}_{\mathcal{Z}} \left( \overbrace{W_k, \dots, W_k}^{M_k}, \dots, \overbrace{W_1, \dots, W_1}^{M_1} \right),$$

taking

$$(3.7) \quad \overbrace{w_k \otimes \cdots \otimes w_k}^{M_k} \otimes \cdots \otimes \overbrace{w_1 \otimes \cdots \otimes w_1}^{M_1}$$

as cyclic vector.

**Lemma 3.2.** *The filtered tensor product (3.6) as  $\mathfrak{sl}_2[t]$ -module and with cyclic vector (3.7) coincides with the filtered tensor product as  $\mathfrak{sl}_3[t]$ -module and with cyclic vector*

$$(3.8) \quad \overbrace{v_k \otimes \cdots \otimes v_k}^{M_k} \otimes \cdots \otimes \overbrace{v_1 \otimes \cdots \otimes v_1}^{M_1}.$$

The proof is given in Appendix, Proposition A.2. Note that the fusion product in the latter sense is the same as the one over the abelian Lie algebra  $\mathfrak{a}[t]$ ,  $\mathfrak{a} = \mathbb{C}e_{13} \oplus \mathbb{C}e_{23}$ . That is,

$$(3.9) \quad W^{(3)}[\lambda] = \overbrace{W_k * \cdots * W_k}^{M_k} * \cdots * \overbrace{W_1 * \cdots * W_1}^{M_1}$$

in the notation of the previous section.

Let  $Y_r$  denote the right ideal of  $U(\mathfrak{sl}_2[t])$  generated by

$$e_{23}[0]^{r+1}, e_{32}[0], \quad x[i] \quad (x \in \mathfrak{sl}_2 \subset \mathfrak{sl}_3, i > 0).$$

As a left  $\mathfrak{sl}_2[t]$ -module we have

$$U(\mathfrak{sl}_2[t])/S(Y_r) \simeq W_r,$$

where  $x[i]$  ( $i > 0$ ) acts as 0 on the right hand side. Introduce the fusion right ideal in the sense of [FJKLM]

$$(3.10) \quad Y_{\lambda} = \overbrace{Y_k \otimes \cdots \otimes Y_k}^{M_k} \otimes \cdots \otimes \overbrace{Y_1 \otimes \cdots \otimes Y_1}^{M_1}(\mathbb{Z}).$$

Our aim is to determine the character of the space of coinvariants

$$(3.11) \quad \text{gr } L_l^{(k)}(\infty)/Y_{\lambda}.$$

Concerning (3.11), the following are known.

**Proposition 3.3.** ([FJKLM], Theorem 2.9) *The dimension of (3.11) is given by the Verlinde rule*

$$\dim L_l^{(k)}(\infty)/Y_{\lambda} = \left( ([0] + \cdots + [k])^{M_k} \cdots ([0] + [1])^{M_1} : [l] \right)_k.$$

From (3.3), the right hand side is written explicitly as

$$(3.12) \quad \sum_{\substack{\mu \subset \lambda \\ |\mu| \equiv l \pmod{2}}} \prod_{i=1}^k \binom{\lambda_i - \mu_{i+1}}{\mu_i - \mu_{i+1}} \cdot K_{l,\mu}^{(k)}(1) = \sum_{\substack{\mu \subset \lambda \\ |\mu| \equiv l \pmod{2}}} F_{\lambda,\mu}(1) K_{l,\mu}^{(k)}(1).$$

**Proposition 3.4.** ([FJKLM], Theorem 3.6) *There is an isomorphism of filtered vector space*

$$(3.13) \quad L_l^{(k)}(\infty)/Y_{\lambda} \simeq \mathcal{F}_{\mathbb{Z}}(\overbrace{W_k, \dots, W_k}^{M_k}, \dots, \overbrace{W_1, \dots, W_1}^{M_1})/S(I_l^{(k)}).$$

On the space  $L_l^{(k)}(\infty)$  we have a grading by  $d = td/dt$  and  $h[0]$ . Define the character of the associated graded space  $L = \text{gr } L_l^{(k)}(\infty)/Y_{\lambda}$  by

$$\text{ch}_{q,z} L = \text{tr}_L(q^d z^{h[0]}).$$

We have also a grading on the fusion product  $W^{(3)}[\lambda]$ . Its character is defined in (2.7), by assigning  $q^i z_1$  (resp.  $q^i z_2$ ) to  $e_{23}[i]$  (resp.  $e_{13}[i]$ ) and 1 to the cyclic vector. Using the action of  $h_{ab}[0] \in \mathfrak{sl}_3$ , we can write the character as

$$\chi^{(3)}[\lambda](q, z_1, z_2) = (z_1 z_2)^{|\lambda|/3} \text{tr}_{W^{(3)}[\lambda]}(q^d z_1^{\bar{\varepsilon}_2} z_2^{\bar{\varepsilon}_1}),$$

where

$$\bar{\varepsilon}_1 = \frac{1}{3}(h_{12}[0] + h_{13}[0]), \quad \bar{\varepsilon}_2 = \frac{1}{3}(h_{21}[0] + h_{23}[0]).$$

**Lemma 3.5.** *In the isomorphism (3.13), the grading by  $h[0]$  in the left hand side translates in the right hand side to the grading by the operator  $(h_{21}[0] + h_{31}[0] + 2|\lambda|)/3$ .*

*Proof.* Consider the cyclic vector  $w_r = v_0 + \cdots + v_r$  of  $W_r$ . As it is explained in (3.20), [FJKLM], the grading by  $h[0]$  counts  $xv_j$  as  $j$  for any  $x \in \mathfrak{sl}_2$ . Denote the natural basis of  $\mathbb{C}^3$  by  $u_i$  ( $i = 1, 2, 3$ ), on which  $e_{jj}[0]$  acts as  $e_{jj}[0]u_i = \delta_{ij}u_i$ . In the identification  $W_r \simeq S^r(\mathbb{C}^3)$ ,  $v_j$  is identified with the symmetrization of  $u_1^{\otimes r-j} \otimes u_3^{\otimes j}$ . Therefore  $h[0]$  acts on  $S^r(\mathbb{C}^3)$  as

$$e_{22}[0] + e_{33}[0] = \frac{1}{3}(h_{21}[0] + h_{31}[0] + 2r).$$

The lemma follows from this.  $\square$

From the lemma we find the following relation

$$\text{ch}_{q,z} \text{gr } L_l^{(k)}(\infty)/Y_\lambda = z^{|\lambda|} \text{ch}_{q,z_1,z_2} W^{(3)}[\lambda]/S(I_l^{(k)})|_{z_1=1, z_2=z^{-1}},$$

where we have used

$$z_1^{\bar{e}_2} z_2^{\bar{e}_1} = z_1^{h_{23}[0]/2} (z_1^{1/2} z_2^{-1})^{-h[0]+2|\lambda|/3}.$$

**3.3. Character formulas.** With the preparations above, let us proceed to the character of a quotient of the fusion product.

**Proposition 3.6.** *We have*

$$\text{ch}_{q,z_1,z_2} W^{(3)}[\lambda]/S(I_l^{(k)}) = \sum_{\substack{\mu \subset \lambda \\ |\mu| \equiv l \pmod{2}}} z_1^{\frac{|\mu|-l}{2}} z_2^{|\lambda|-|\mu|} F_{\lambda,\mu}(q) K_{l,\mu}^{(k)}(q),$$

where  $F_{\lambda,\mu}(q)$  is defined in (2.49).

*Proof.* By Proposition 3.3 and 3.4, (3.12) gives a lower bound to the dimension  $\dim W^{(3)}[\lambda]/S(I_l^{(k)})$ .

On the other hand, from Proposition 2.10 we have a filtration of  $W^{(3)}[\lambda]$  by submodules with composition factors of the form  $W^{(2)}[\mu]$ . The operators  $h_{23}[0] + l$ ,  $e_{23}[0]$  and  $e_{23}[1]^{k-l+1}$  leave the filtration invariant. Replacing the character of the quotient by that of the quotient of the associated graded space, we obtain an upper estimate

$$(3.14) \quad \begin{aligned} & \text{ch}_{q,z_1,z_2} W^{(3)}[\lambda]/S(I_l^{(k)}) \\ & \leq \sum_{\substack{\mu \subset \lambda \\ |\mu| \equiv l \pmod{2}}} z_2^{|\lambda|-|\mu|} F_{\lambda,\mu}(q) \text{ch}_{q,z_1} W^{(2)}[\mu]/S(I_l^{(k)}). \end{aligned}$$

Proposition 3.1 states that up to a power of  $z_1$  the last factor is given by  $K_{l,\mu}^{(k)}(q)$ . Setting  $q = z_1 = z_2 = 1$ , the right hand side of (3.14) becomes the lower bound (3.12). This in turn implies that the equality holds in each step, and the proof is over.  $\square$



Writing  $F_{\lambda\mu}(q)$  explicitly, we obtain the following as an immediate consequence of Proposition 3.1 and 3.4.

**Theorem 3.7.** *We have a fermionic formula for the character of the space of coinvariants*

$$(3.15) \quad \text{ch}_{q,z} \text{gr } L_l^{(k)}(\infty)/Y_\lambda$$

$$= \sum_{\substack{\mu \subset \lambda \\ |\mu| \equiv l \pmod{2}}} z^{|\mu|} q^{\sum_{i=1}^{k-1} \mu_{i+1}(\lambda_i - \mu_i)} \prod_{i=1}^k \begin{bmatrix} \lambda_i - \mu_{i+1} \\ \mu_i - \mu_{i+1} \end{bmatrix} \cdot K_{l,\mu}^{(k)}(q).$$

Since  $W^{(3)}[\lambda]$  has an  $\mathfrak{sl}_2$ -structure, its character is a linear combination of  $\mathfrak{sl}_2$ -characters  $\chi_j(z) = z^{-j/2} + \dots + z^{j/2-1} + z^{j/2}$ . Let us define  $X_{j\lambda}(q, z)$  as the coefficient of the expansion

$$\chi^{(3)}[\lambda](q, z_1, z_2) = z_2^{|\lambda|} \sum_{j \geq 0} \chi_j(z_1) X_{j\lambda}(q, z_1^{1/2} z_2^{-1}).$$

Then we have the following alternating sum formula. When specialized to the case  $\mathbf{M} = (0, \dots, 0, M)$ , it settles the ‘main conjecture’ of [FL] (Conjecture 3.5) for  $\mathfrak{g} = \mathfrak{sl}_2$ .

**Theorem 3.8.**

$$(3.16) \text{ch}_{q,z} \text{gr } L_l^{(k)}(\infty)/Y_\lambda = \sum_{i \geq 0} q^{(k+2)i^2 + (l+1)i} X_{2(k+2)i+l,\lambda}(q, z)$$

$$- \sum_{i > 0} q^{(k+2)i^2 - (l+1)i} X_{2(k+2)i-l-2,\lambda}(q, z).$$

*Proof.* We have from (2.50) and (2.52)

$$(3.17) \quad \chi^{(3)}[\lambda](q, z_1, z_2) = \sum_{\mu \subset \lambda} z_2^{|\lambda| - |\mu|} F_{\lambda\mu}(q) \chi^{(2)}[\mu](q, z_1),$$

$$(3.18) \quad \chi^{(2)}[\mu](q, z_1) = \sum_{0 \leq j \leq |\mu|} z_1^{(|\mu| - j)/2} \tilde{S}_{\sigma(j), \mu}(q),$$

where we have set  $\sigma(j) = (\frac{|\mu|+j}{2}, \frac{|\mu|-j}{2})$ . Let  $K_{j,\mu}(q)$  denote the non-restricted Kostka polynomial for  $\mathfrak{sl}_2$ , which is related to the  $q$ -supernomial coefficients  $\tilde{S}_{\sigma(j), \mu}(q)$  via  $\chi^{(2)}[\mu](q, z_1)$  as

$$\chi^{(2)}[\mu](q, z_1) = \sum_{0 \leq j \leq |\mu|} z_1^{|\mu|/2} \chi_j(z_1) K_{j,\mu}(q).$$

From this and (3.17), (3.18), we find an expression

$$X_{j\lambda}(q, z) = \sum_{|\mu|=j} z^{|\mu|} F_{\lambda\mu}(q) K_{j,\mu}(q).$$

The statement of Theorem now follows from Theorem 3.7 together with the known formula (see [SS], eq.(6.8))

$$\begin{aligned} K_{l,\mu}^{(k)}(q) &= \sum_{i \geq 0} q^{(k+2)i^2 + (l+1)i} K_{2(k+2)i+l,\mu}(q) \\ &\quad - \sum_{i > 0} q^{(k+2)i^2 - (l+1)i} K_{2(k+2)i-l-2,\mu}(q). \end{aligned}$$

□

#### APPENDIX A. A LEMMA ON CHANGING CYCLIC VECTORS

In this appendix we prove one of the steps used in the main text in a more general setting, i.e., to show the equality of certain filtered tensor products of  $\mathfrak{sl}_{n-1}[t]$  and  $\mathfrak{sl}_n[t]$  representations. In this paper, we use only the case of  $(\mathfrak{sl}_2, \mathfrak{sl}_3)$  pair and for symmetric tensor representations.

Since the essential part of the argument is contained in the simplest case  $n = 2$ , we first formulate it and give a proof to that particular case. Consider the subalgebra  $\mathbb{C}h \otimes \mathbb{C}[t] \subset \mathfrak{sl}_2[t]$ .

**Proposition A.1.** *Let  $v_a$  ( $1 \leq a \leq N$ ) be the lowest weight vectors of finite dimensional irreducible representations of  $\mathfrak{sl}_2$ . We denote by  $U_a$  the representation generated by  $v_a$ . Let  $\mathcal{Z} = (z_1, \dots, z_N)$  be (not necessarily distinct) complex numbers. Consider the tensor product of the evaluation representations  $U_a$  ( $1 \leq a \leq N$ ) of  $\mathfrak{sl}_2[t]$ , for which the evaluation parameter is set to  $z_a$ . We denote this tensor product by  $W(\mathcal{Z})$ .*

*Let  $w_0 = v_1 \otimes \dots \otimes v_N$ , and consider the subspace  $G^i$  of  $W(\mathcal{Z})$  spanned by the vectors  $x_1[i_1] \dots x_l[i_l]w_0$  where  $l \geq 0$ ,  $x_1, \dots, x_l \in \mathfrak{sl}_2$  and  $\sum_a i_a \leq i$ .*

*Let*

$$w = \left( \sum_i \frac{e[0]^i}{i!} v_1 \right) \otimes \dots \otimes \left( \sum_i \frac{e[0]^i}{i!} v_N \right),$$

*and let  $F^i$  be the subspace of  $W(\mathcal{Z})$  spanned by the vectors  $h[i_1] \dots h[i_l]w$  where  $l \geq 0$  and  $\sum_a i_a \leq i$ .*

*Under this setting, the equality  $F^i = G^i$  holds for all  $i \geq 0$ .*

*Proof.* Note that

$$w = \sum_i \frac{e[0]^i}{i!} w_0.$$

Since the  $h[0]$ -weights of the vectors  $e[0]^i w_0$  ( $i \geq 0$ ) are distinct, they all belong to  $F^0$ . Since  $\sum_i \mathbb{C}e[0]^i w_0$  is  $h \otimes t^0$ -invariant, we have  $F^0 = \sum_i \mathbb{C}e[0]^i w_0$ . Since  $f[0]w_0, h[0]w_0 \in \mathbb{C}w_0$ ,  $F^0$  is  $\mathfrak{sl}_2 \otimes t^0$  invariant, and therefore,  $F^0 = G^0$ .

Now, we will show that  $(\mathfrak{sl}_2 \otimes t^j)F^i \subset F^{i+j}$  for all  $i, j \geq 0$ . From this follows  $F^i = G^i$  for all  $i$ . Note that

$$F^i = \sum_{m \geq 0} \sum_{i_1 + \dots + i_K \leq i} \mathbb{C}h[i_1] \dots h[i_K] e[0]^m w_0.$$

First, we show that

$$e[j]h[i_1] \cdots h[i_K]e[0]^m w_0 \in F^{i_1 + \cdots + i_K + j}$$

by induction on  $K$ . For  $K = 0$ , we have

$$\begin{aligned} e[j]e[0]^m w_0 &= e[0]^m e[j]w_0 \\ &= \frac{1}{2}e[0]^m [h[j], e[0]]w_0. \end{aligned}$$

We use  $a \equiv_i b$  meaning  $a - b \in F^i$ . Since  $h[j]w_0 \in \mathbf{C}w_0$  and  $h[j]e[0]^{m+1}w_0 \in F^j$ , we have

$$\begin{aligned} e[j]e[0]^m w_0 &\equiv_j \frac{1}{2}e[0]^m h[j]e_0 w_0 \\ &\equiv_j \frac{1}{2}[e[0]^m, h[j]]e[0]w_0 \\ &= -me[j]e[0]^m w_0. \end{aligned}$$

From this follows that  $e[j]e[0]^m w_0 \in F^j$ .

For  $K \geq 1$ , we have

$$e[j]h[i_1] \cdots h[i_K]e[0]^m w_0 = [e[j], h[i_1]] \cdots h[i_K]w_0 + h[i_1]e[j] \cdots h[i_K]w_0.$$

Using induction hypothesis, we see that this element belongs to  $F^{i_1 + \cdots + i_K + j}$ .  $\square$

Now, we proceed to the case of  $(\mathfrak{sl}_{n-1}, \mathfrak{sl}_n)$  pair.

**Proposition A.2.** *Let  $v_a$  ( $1 \leq a \leq N$ ) be the lowest weight vectors of finite dimensional irreducible representations of  $\mathfrak{sl}_n$  ( $n \geq 3$ ). We denote by  $U_a$  the representation generated by  $v_a$ . Let  $\mathcal{Z} = (z_1, \dots, z_N)$  be (not necessarily distinct) complex numbers. Consider the tensor product of the evaluation representations  $U_a$  ( $1 \leq a \leq N$ ) of  $\mathfrak{sl}_n[t]$ , for which the evaluation parameter is set to  $z_a$ . We denote this tensor product by  $W(\mathcal{Z})$ .*

*Let  $w_0 = v_1 \otimes \cdots \otimes v_N$ , and consider the subspace  $G^i$  of  $W(\mathcal{Z})$  spanned by the vectors  $x_1[i_1] \cdots x_l[i_l]w_0$  where  $l \geq 0$ ,  $x_1, \dots, x_l \in \mathfrak{sl}_n$  and  $\sum_a i_a \leq i$ .*

*Let*

$$w = \left( \sum_i \frac{e_{1n}^i}{i!} v_1 \right) \otimes \cdots \otimes \left( \sum_i \frac{e_{1n}^i}{i!} v_N \right),$$

*and let  $F^i$  be the subspace of  $W(\mathcal{Z})$  spanned by the vectors  $x_1[i_1] \cdots x_l[i_l]w$  where  $l \geq 0$ ,  $x_1, \dots, x_l \in \mathfrak{sl}_{n-1}$  and  $\sum_a i_a \leq i$ .*

*Under this setting, the equality  $F^i = G^i$  holds for all  $i \geq 0$ .*

*Proof. First Step:* We show that  $F^0$  is spanned by the vectors

$$(A.1) \quad e_{a_1 b_1}[0] \cdots e_{a_J b_J}[0] \prod_{2 \leq a \leq n} e_{1a}[0]^{m_a} w_0,$$

where  $J \geq 0$ ,  $2 \leq a_i < b_i \leq n$  for  $1 \leq i \leq J$ , and  $m_a \geq 0$  for  $2 \leq a \leq n$ .

Proceeding as in the proof of Proposition A.1, we see that  $F^0$  contains the vectors  $e_{1n}[0]^m w_0$  ( $m \geq 0$ ). Then, by using  $[e_{na}[0], e_{1b}[0]] = -\delta_{n,b} e_{1a}[0]$

and  $[e_{1a}[0], e_{1b}[0]] = 0$  for  $2 \leq a \leq n-1$  and  $2 \leq b \leq n$ , we see that  $F^0$  contains the vectors  $\prod_{2 \leq a \leq n} e_{1a}[0]^{m_a} w_0$  for  $m_2, \dots, m_n \geq 0$ .

Now, by using the PBW theorem, it is straightforward to see that  $F^0$  is spanned by the vectors (A.1).

*Second Step:* We show that  $F^0 = G^0$ . It is enough to show that  $F^0$  is invariant by the actions of  $e_{1a}[0]$  and  $e_{a1}[0]$  for  $2 \leq a \leq n$ . The invariance for  $e_{1a}[0]$  follows from  $[e_{1a}[0], e_{bc}[0]] = \delta_{ab} e_{1c}[0]$  for  $2 \leq a \leq n$  and  $2 \leq b < c \leq n$ . The invariance for  $e_{a1}[0]$  follows from  $e_{a1}[0]w_0 = 0$ ,  $[e_{a1}[0], e_{bc}[0]] = -\delta_{ac} e_{b1}[0]$  for  $2 \leq a \leq n$  and  $2 \leq b < c \leq n$ , and

$$[e_{a1}[0], e_{1b}[0]] = \begin{cases} -h_{1a}[0] & \text{if } a = b; \\ e_{ab}[0] & \text{if } a \neq b, \end{cases}$$

for  $2 \leq a, b \leq n$ .

*Third Step:* We show that  $F^j$  contains the vectors

$$(A.2) \quad e_{a_1 b_1}[j_1] \cdots e_{a_J b_J}[j_J] \prod_{i=1}^K h_{c_i}[k_i] \prod_{i=1}^L e_{1d_i}[l_i] e_{1n}[0]^m w_0,$$

where  $J, K, L, j_i, l_i, m \geq 0$ ,  $k_i > 0$ ,  $2 \leq a_i < b_i \leq n$ ,  $2 \leq c_i, d_i \leq n-1$ , and

$$\sum_{i=1}^J j_i + \sum_{i=1}^K k_i + \sum_{i=1}^L l_i \leq j.$$

We have shown that  $F^0$  contains the vectors  $\prod_{a=2}^n e_{1a}[0]^{m_a} w_0$ . Applying  $e_{32}[l_i^{(2)}]$  ( $1 \leq i \leq m'_2 \leq m_3; l_i^{(2)} > 0$ ), we get

$$\prod_{a=4}^n e_{1a}[0]^{m_a} e_{13}[0]^{m_3-m'_2} \prod_{i=1}^{m'_2} e_{12}[l_i^{(2)}] e_{12}[0]^{m_2} w_0 \in F^{\sum_{i=1}^{m'_2} l_i^{(2)}}$$

Repeating this procedure successively for  $e_{43}, \dots, e_{n,n-1}$ , we obtain the vectors  $\prod_{i=1}^L e_{1d_i}[l_i] e_{1n}[0]^m w_0 \in F^{\sum_{i=1}^L l_i}$ . Therefore,  $F^j$  contains the vectors (A.2). We denote by  $H^j$  the linear span of these vectors.

*Last Step:* We show that

$$(\mathfrak{sl}_n \otimes t^i) H^j \subset H^{i+j}.$$

Since  $H^0 = F^0 = G^0$ , it then follows that  $H^j = F^j = G^j$ . The only non-trivial calculation is to show that  $e_{1n}[i] H^j \subset H^{i+j}$ . This can be shown similarly as in the proof of Proposition A.1.  $\square$

Let us explain these statements in more invariant terms. To define a lowest vector of an irreducible representation it is enough to choose a Borel subalgebra  $\mathfrak{b} \subset \mathfrak{sl}_n$ . The set of all possible Borel subalgebras inherits the natural topology of a homogeneous space, namely they form the flag manifold.

Now take the standard inclusion  $\mathfrak{sl}_{n-1} \subset \mathfrak{sl}_n$  and choose a Borel subalgebra  $\mathfrak{b} \subset \mathfrak{sl}_n$ . Then for any irreducible representation  $U$  of  $\mathfrak{sl}_n$  by  $v_U^{\mathfrak{b}}$  denote

the corresponding lowest weight vector and by  $\overline{U}$  denote the restriction of  $U$  to  $\mathfrak{sl}_{n-1}$ .

**Theorem A.3.** *Consider a set of irreducible representations  $U_1, \dots, U_N$  of  $\mathfrak{sl}_n$  and a set of distinct points  $\mathcal{Z} = (z_1, \dots, z_n)$ . Choose a generic Borel subalgebra  $\mathfrak{b} \subset \mathfrak{sl}_n$ . Then vectors  $v_{U_a}^{\mathfrak{b}}$  are cyclic in  $\overline{U}_a$  as well as in  $U_a$  and we have an isomorphism of filtered spaces*

$$(A.3) \quad \mathcal{F}_{\mathcal{Z}}(U_1, \dots, U_N) \cong \mathcal{F}_{\mathcal{Z}}(\overline{U}_1, \dots, \overline{U}_N).$$

*Proof.* Note, that (A.3) is satisfied for an algebraically open set of Borel subalgebras. So it is enough to show it for a certain subalgebra.

Let  $\mathfrak{b}_0$  be the standard Borel subalgebra of upper-triangular matrices. Then Proposition A.2 implies (A.3) for

$$\mathfrak{b} = \exp(e_{1n})\mathfrak{b}_0 \exp(-e_{1n}).$$

□

*Acknowledgments.* This work is partially supported by the Grant-in-Aid for Scientific Research (B2) no.12440039, no.14340040 and (A1) no.13304010, Japan Society for the Promotion of Science. BF is partially supported by grants RFBR 02-01-01015 and INTAS-00-00055. The work of SL is partially supported by the grant RFBR-01-01-00546. The last stage of this work was carried out while the authors were visiting Mathematical Sciences Research Institute, Berkeley, March 2002.

## REFERENCES

- [FF] B. L. Feigin and E. Feigin,  $q$ -characters of the tensor products in  $\mathfrak{sl}_2$ -case, math.QA/0201111 (2002).
- [FFu] B. L. Feigin and D. B. Fuchs, Cohomology of some nilpotent subalgebras of the Virasoro and Kac-Moody algebras, *Geometry and Physics*, Essays in honor of I.M.Gelfand on the occasion of his 75th birthday, eds. S.Gindikin and I.M.Singer, 209–235, North Holland, (1991)
- [FJKLM] B. Feigin, M. Jimbo, R. Kedem, S. Loktev and T. Miwa, Spaces of Coinvariants and Fusion Product I. From Equivalence Theorem to Kostka Polynomials. math.QA/
- [FKLMM] B. Feigin, R. Kedem, S. Loktev, T. Miwa and E. Mukhin, Combinatorics of the  $\widehat{\mathfrak{sl}}_2$  spaces of coinvariants, *Transformation Groups*, **6** (2001) 25–52; Combinatorics of the  $\widehat{\mathfrak{sl}}_2$  spaces of coinvariants: Loop Heisenberg modules and recursion, math.QA/0009198; Combinatorics of the  $\widehat{\mathfrak{sl}}_2$  spaces of coinvariants: Dual functional realization and recursion, math.QA/0012190.
- [FL] B. L. Feigin and S. Loktev, On generalized Kostka polynomials and quantum Verlinde rule, math.QA/9812093, *Amer. Math. Sci. Transl.* **194** (1999) 61–79.
- [FS] B. L. Feigin and A. V. Stoyanovsky, Functional models for representations of current algebras and semi-infinite Schubert cells, *Funct. Anal. and Its Appl.* **28** (1993) 55–72.
- [HKKOTY] G. Hatayama, A. N. Kirillov, A. Kuniba, M. Okado, T. Takagi and Y. Yamada, Character formulae of  $\widehat{\mathfrak{sl}}_n$ -modules and inhomogeneous paths, math.QA/9802085, *Nucl. Phys. B* **536** (1999) 575–616.

- [KKMM] R. Kedem, T. R. Klassen, B. M. McCoy and E. Melzer, Fermionic sum representations for conformal field theory characters, *Phys. Lett. B* **307** (1993) 68–76.
- [KKMOTU] A. Kuniba, K. M. Misra, M. Okado, T. Takagi and J. Uchiyama, Characters of Demazure modules and solvable lattice models, q-alg.9707004, *Nucl. Phys. B* **536** (1999) 575–616.
- [KM] R. Kedem and B. M. McCoy, Construction of modular branching functions from Bethe’s equations in the 3-state Potts chain, *J. Stat. Phys.* **71** (1993) 875–901.
- [KR] A.N. Kirillov and N.Yu Reshetikhin, The Bethe Ansatz and the combinatorics of Young tableaux, *J. Soviet Math.* **41** (1988) 925–955.
- [LS] A. Lascoux and M.P. Schützenberger, Sur une conjecture de H. O. Foulkes, *C. R. Acad. Sci. Paris Ser. A-B* **286** (1978) A323–A324.
- [NY] A. Nakayashiki and Y. Yamada, Kostka polynomials and energy functions in solvable lattice models, *Selecta Math.* **3** (1997) 547–599.
- [S] A. Schilling,  $q$ -Supernomial coefficients: From riggings to ribbons, math.CO/0107214, to appear in ‘MathPhys Odessey 2001’, Birkhäuser.
- [SS] A. Schilling and M. Shimozono, Fermionic formulas for level-restricted generalized Kostka polynomials and coset branching functions, math.QA/0001114, *Commun. Math. Phys.* **220** (2001) 105–164.
- [SW1] A. Schilling and S. O. Warnaar, Supernomial coefficients, polynomial identities and  $q$ -series, math.QA/9701007, *Ramanujan J.* **2** (1998) 459–494.
- [SW2] A. Schilling and S. O. Warnaar, Inhomogeneous lattice paths, generalized Kostka polynomials and  $A_{n-1}$  supernomials, math.QA/9802111, *Commun. Math. Phys.* **202** (1999) 359–401.

BF: LANDAU INSTITUTE FOR THEORETICAL PHYSICS, CHERNOGOLOVKA, 142432, RUSSIA

*E-mail address:* feigin@feigin.mccme.ru

MJ: GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, TOKYO 153-8914, JAPAN

*E-mail address:* jimbomic@ms.u-tokyo.ac.jp

RK: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, 1409 W. GREEN ST. URBANA, IL 61801, USA

*E-mail address:* rinat@math.uiuc.edu

SL: INSTITUTE FOR THEORETICAL AND EXPERIMENTAL PHYSICS, B. CHEREMUSHKINSKAYA, 25, MOSCOW 117259, RUSSIA

INDEPENDENT UNIVERSITY OF MOSCOW, B. VLASIEVSKY PER, 11, MOSCOW 121002, RUSSIA

*E-mail address:* loktev@mccme.ru

TM: DIVISION OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, KYOTO UNIVERSITY, KYOTO 606-8502 JAPAN

*E-mail address:* tetsuji@kusm.kyoto-u.ac.jp